

HSD17B1 Antibody



PACO55550

Product Information

Size:

50ug

Reactivity:

Human

Source:

Rabbit

Isotype:

IgG

Applications:

ELISA, IHC, IF

Recommended dilutions:

ELISA:1:2000-1:10000, IHC:1:20-1:200,
IF:1:20-1:200

Protein Background:

Favors the reduction of estrogens and androgens. Also has 20-alpha-HSD activity. Uses preferentially NADH.

Gene ID:

HSD17B1

Uniprot

P14061

Synonyms:

Estradiol 17-beta-dehydrogenase 1 (EC 1.1.1.62) (17-beta-hydroxysteroid dehydrogenase type 1) (17-beta-HSD 1) (20 alpha-hydroxysteroid dehydrogenase) (20-alpha-HSD) (E2DH) (Placental 17-beta-hydroxysteroid dehydrogenase) (Short chain dehydrogenase/reductase family 28C member 1), HSD17B1, E17KSR EDH17B1 EDH17B2 EDHB17 SDR28C1


Immunogen:

Recombinant Human Estradiol 17-beta-dehydrogenase 1 protein (268-328AA).


Storage:

Preservative: 0.03% Proclin 300. Constituents: 50% Glycerol, 0.01M PBS, pH 7.4


Product Images



IHC image of PACO55550 diluted at 1:100 and staining in paraffin-embedded human prostate cancer performed on a Leica Bond™ system. After dewaxing and hydration, antigen retrieval was mediated by high pressure in a citrate buffer (pH 6.0). Section was blocked with 10% normal goat serum 30min at RT. Then primary antibody (1% BSA) was incubated at 4°C overnight. The primary is detected by a biotinylated secondary antibody and visualized using an HRP conjugated SP system.



Immunofluorescence staining of HeLa cells with PACO55550 at 1:33, counter-stained with DAPI. The cells were fixed in 4% formaldehyde, permeabilized using 0.2% Triton X-100 and blocked in 10% normal Goat Serum. The cells were then incubated with the antibody overnight at 4°C. The secondary antibody was Alexa Fluor 488-conjugated AffiniPure Goat Anti-Rabbit IgG(H+L).



IHC image of PACO55550 diluted at 1:100 and staining in paraffin-embedded human colon cancer performed on a Leica Bond™ system. After dewaxing and hydration, antigen retrieval was mediated by high pressure in a citrate buffer (pH 6.0). Section was blocked with 10% normal goat serum 30min at RT. Then primary antibody (1% BSA) was incubated at 4°C overnight. The primary is detected by a biotinylated secondary antibody and visualized using an HRP conjugated SP system.