## **UBA52 Antibody**



## PACO55818

**Product Information** 

Size: Protein Background:

50ug

Reactivity: Gene ID:

Human UBA52

Source: Uniprot

Rabbit P62987

Isotype: Synonyms:

lgG Ubiquitin-60S ribosomal protein L40 (CEP52) (Ubiquitin A-52 residue ribosomal protein

fusion product 1) [Cleaved into: Ubiquitin; 60S ribosomal protein L40 (Large ribosomal

**Applications:** subunit protein eL40)], UBA52, UBCEP2

ELISA, IHC, IF Immunogen:

**Recommended dilutions:** Recombinant Human Ubiquitin-60S ribosomal protein L40 protein (1-128AA).


ELISA:1:2000-1:10000, IHC:1:200-1:500,


IF:1:50-1:200

Storage:

Preservative: 0.03% Proclin 300. Constituents: 50% Glycerol, 0.01M PBS, pH 7.4


## **Product Images**





IHC image of PACO55818 diluted at 1:200 and staining in paraffinembedded human testis tissue performed on a Leica BondTM system. After dewaxing and hydration, antigen retrieval was mediated by high pressure in a citrate buffer (pH 6.0). Section was blocked with 10% normal goat serum 30min at RT. Then primary antibody (1% BSA) was incubated at 4°C overnight. The primary is detected by a biotinylated secondary antibody and visualized using an HRP conjugated SP system.

Immunofluorescence staining of MCF-7 cells with PACO55818 at 1:66, counter-stained with DAPI. The cells were fixed in 4% formaldehyde, permeabilized using 0.2% Triton X-100 and blocked in 10% normal Goat Serum. The cells were then incubated with the antibody overnight at 4°C. The secondary antibody was Alexa Fluor 488-congugated AffiniPure Goat Anti-Rabbit IgG(H+L).



IHC image of PACO55818 diluted at 1:200 and staining in paraffinembedded human brain tissue performed on a Leica BondTM system. After dewaxing and hydration, antigen retrieval was mediated by high pressure in a citrate buffer (pH 6.0). Section was blocked with 10% normal goat serum 30min at RT. Then primary antibody (1% BSA) was incubated at 4°C overnight. The primary is detected by a biotinylated secondary antibody and visualized using an HRP conjugated SP system.