## HIST1H3A (Ab-36) Antibody



## PACO56506

Reactivity:

Human

Rabbit

## **Product Information**

Size: **Protein Background:** 

50ul Core component of nucleosome. Nucleosomes wrap and compact DNA into chromatin,

> limiting DNA accessibility to the cellular machineries which require DNA as a template. Histones thereby play a central role in transcription regulation, DNA repair, DNA

replication and chromosomal stability. DNA accessibility is regulated via a complex set

of post-translational modifications of histones, also called histone code, and

Source: nucleosome remodeling.

Gene ID:

HIST1H3A Isotype:

lgG Uniprot

P68431 **Applications:** 

ELISA, IHC, IF Synonyms:

**Recommended dilutions:** 

ELISA:1:2000-1:10000, IHC:1:20-1:200, IF:1:50-1:200

Histone H3.1 (Histone H3/a) (Histone H3/b) (Histone H3/c) (Histone H3/d) (Histone H3/f) (Histone H3/h) (Histone H3/i) (Histone H3/j) (Histone H3/k) (Histone H3/l), HIST1H3A; HIST1H3B; HIST1H3C; HIST1H3D; HIST1H3E; HIST1H3F; HIST1H3G; HIST1H3H; HIST1H3I; HIST1H3J, H3FA; H3FL; H3FC; H3FB; H3FD; H3FI; H3FH; H3FK; H3FF; H3FJ


Immunogen:


Peptide sequence around site of Lys (36) derived from Human Histone H3.1.


Storage:

Preservative: 0.03% Proclin 300. Constituents: 50% Glycerol, 0.01M PBS, pH 7.4


## **Product Images**







Immunofluorescent analysis of Hela cells using PACO56506 at dilution of 1:100 and Cy3-congugated Goat Anti-Rabbit IgG.



Immunohistochemistry of paraffin-embedded human lung cancer using PACO56506 at dilution of 1:100.