# HIST1H3A (Ab-9) Antibody



#### PACO56521

Reactivity:

Human

Source:

#### **Product Information**

Size: Protein Background:

50ul Core component of nucleosome. Nucleosomes wrap and compact DNA into chromatin, limiting DNA accessibility to the cellular machineries which require DNA as a template.

Histones thereby play a central role in transcription regulation, DNA repair, DNA

replication and chromosomal stability. DNA accessibility is regulated via a complex set

of post-translational modifications of histones, also called histone code, and

nucleosome remodeling.

Rabbit Gene ID:

Isotype: HIST1H3A

lgG Uniprot

Applications: P68431

ELISA, WB, IHC, IF Synonyms:

Recommended dilutions:

ELISA:1:2000-1:10000, WB:1:100-1:1000, IHC:1:20-1:200, IF:1:50-1:200

Histone H3.1 (Histone H3/a) (Histone H3/b) (Histone H3/c) (Histone H3/d) (Histone H3/f) (Histone H3/f) (Histone H3/f) (Histone H3/f) (Histone H3/f) (Histone H3/f) (Histone H3/f), HIST1H3A; HIST1H3B; HIST1H3C; HIST1H3D; HIST1H3E; HIST1H3F; HIST1H3G; HIST1H3H; HIST1H3I; HIST1H3J, H3FA; H3FL; H3FC; H3FB; H3FD; H3FI; H3FH; H3FK; H3FF; H3FJ


## Immunogen:

Peptide sequence around site of Lys (9) derived from Human Histone H3.1.


### Storage:

Preservative: 0.03% Proclin 300. Constituents: 50% Glycerol, 0.01M PBS, pH 7.4


## **Product Images**



Western Blot. Positive WB detected in: Hela cell acid extracts. All lanes: HIST1H3A antibody at  $2\mu g/ml$ . Secondary. Goat polyclonal to rabbit lgG at 1/50000 dilution. Predicted band size: 16 kDa. Observed band size: 16 kDa.



Immunofluorescent analysis of Hela cells using PACO56521 at dilution of 1:100 and Cy3-congugated Goat Anti-Rabbit IgG.



Immunohistochemistry of paraffin-embedded human lung cancer using PACO56521 at dilution of 1:100.