

PACO57264

---

## Product Information

**Size:**

50ug

**Reactivity:**

Human

**Source:**

Rabbit

**Isotype:**

IgG

**Applications:**

ELISA, IHC

**Recommended dilutions:**

ELISA:1:2000-1:10000, IHC:1:200-1:500

**Protein Background:**

Supplies decaprenyl diphosphate, the precursor for the side chain of the isoprenoid quinones ubiquinone-10.

**Gene ID:**

PDSS2

**Uniprot**

Q86YH6

**Synonyms:**

Decaprenyl-diphosphate synthase subunit 2 (EC 2.5.1.91) (All-trans-decaprenyl-diphosphate synthase subunit 2) (Candidate tumor suppressor protein) (Decaprenyl pyrophosphate synthase subunit 2), PDSS2, C6orf210 DLP1

**Immunogen:**


Recombinant Human Decaprenyl-diphosphate synthase subunit 2 protein (292-387AA).

**Storage:**


Preservative: 0.03% Proclin 300. Constituents: 50% Glycerol, 0.01M PBS, pH 7.4

## Product Images

---



IHC image of PACO57264 diluted at 1:200 and staining in paraffin-embedded human liver cancer performed on a Leica Bond™ system. After dewaxing and hydration, antigen retrieval was mediated by high pressure in a citrate buffer (pH 6.0). Section was blocked with 10% normal goat serum 30min at RT. Then primary antibody (1% BSA) was incubated at 4°C overnight. The primary is detected by a biotinylated secondary antibody and visualized using an HRP conjugated SP system.



IHC image of PACO57264 diluted at 1:200 and staining in paraffin-embedded human kidney tissue performed on a Leica Bond™ system. After dewaxing and hydration, antigen retrieval was mediated by high pressure in a citrate buffer (pH 6.0). Section was blocked with 10% normal goat serum 30min at RT. Then primary antibody (1% BSA) was incubated at 4°C overnight. The primary is detected by a biotinylated secondary antibody and visualized using an HRP conjugated SP system.