

TRIM8 Antibody



PACO57480

Product Information

Size:

50ug

Reactivity:

Human

Source:

Rabbit

Isotype:

IgG

Applications:

ELISA, IHC, IF

Recommended dilutions:

ELISA:1:2000-1:10000, IHC:1:200-1:500,
IF:1:50-1:200

Protein Background:

Probable E3 ubiquitin-protein ligase which may promote proteasomal degradation of SOCS1.

Gene ID:

TRIM8

Uniprot

Q9BZR9

Synonyms:

E3 ubiquitin-protein ligase TRIM8 (EC 2.3.2.27) (Glioblastoma-expressed RING finger protein) (RING finger protein 27) (RING-type E3 ubiquitin transferase TRIM8) (Tripartite motif-containing protein 8), TRIM8, GERP RNF27


Immunogen:

Recombinant Human Probable E3 ubiquitin-protein ligase TRIM8 protein (367-517AA).


Storage:

Preservative: 0.03% Proclin 300. Constituents: 50% Glycerol, 0.01M PBS, pH 7.4

Product Images



Immunofluorescence staining of HepG2 cells with PACO57480 at 1:133, counter-stained with DAPI. The cells were fixed in 4% formaldehyde, permeabilized using 0.2% Triton X-100 and blocked in 10% normal Goat Serum. The cells were then incubated with the antibody overnight at 4°C. The secondary antibody was Alexa Fluor 488-conjugated AffiniPure Goat Anti-Rabbit IgG(H+L).



IHC image of PACO57480 diluted at 1:400 and staining in paraffin-embedded human kidney tissue performed on a Leica Bond™ system. After dewaxing and hydration, antigen retrieval was mediated by high pressure in a citrate buffer (pH 6.0). Section was blocked with 10% normal goat serum 30min at RT. Then primary antibody (1% BSA) was incubated at 4°C overnight. The primary is detected by a biotinylated secondary antibody and visualized using an HRP conjugated SP system.