
PNPO Antibody

PACO57556



Product Information	
Size:	Protein Background:
50ug	Catalyzes the oxidation of either pyridoxine 5'-phosphate (PNP) or pyridoxamine 5'-
Reactivity:	phosphate (PMP) into pyridoxal 5'-phosphate (PLP).
Human	Gene ID:
Source:	PNPO
	Uniprot
Rabbit	Q9NVS9
lsotype:	
lgG	Synonyms:
Applications:	Pyridoxine-5'-phosphate oxidase (EC 1.4.3.5) (Pyridoxamine-phosphate oxidase), PNPO
	Immunogen:
ELISA, IHC, IF	Recombinant Human Pyridoxine-5'-phosphate oxidase protein (114-207AA).
Recommended dilutions:	
ELISA:1:2000-1:10000, IHC:1:500-1:1000,	Storage:
IF:1:200-1:500	Preservative: 0.03% Proclin 300. Constituents: 50% Glycerol, 0.01M PBS, pH 7.4



IHC image of PACO57556 diluted at 1:600 and staining in paraffinembedded human kidney tissue performed on a Leica BondTM system. After dewaxing and hydration, antigen retrieval was mediated by high pressure in a citrate buffer (pH 6.0). Section was blocked with 10% normal goat serum 30min at RT. Then primary antibody (1% BSA) was incubated at 4°C overnight. The primary is detected by a biotinylated secondary antibody and visualized using an HRP conjugated SP system.

Immunofluorescence staining of Hela cells with PACO57556 at 1:200, counter-stained with DAPI. The cells were fixed in 4% formaldehyde, permeabilized using 0.2% Triton X-100 and blocked in 10% normal Goat Serum. The cells were then incubated with the antibody overnight at 4°C. The secondary antibody was Alexa Fluor 488-congugated AffiniPure Goat Anti-Rabbit IgG(H+L).

IHC image of PACO57556 diluted at 1:600 and staining in paraffinembedded human endometrial cancer performed on a Leica BondTM system. After dewaxing and hydration, antigen retrieval was mediated by high pressure in a citrate buffer (pH 6.0). Section was blocked with 10% normal goat serum 30min at RT. Then primary antibody (1% BSA) was incubated at 4°C overnight. The primary is detected by a biotinylated secondary antibody and visualized using an HRP conjugated SP system.