## **UBIAD1** Antibody, FITC conjugated



## PACO58070

Reactivity:

Human

Isotype:

## **Product Information**

Size: **Protein Background:** 

50ug Prenyltransferase that mediates the formation of menaquinone-4 (MK-4) and coenzyme Q10. MK-4 is a vitamin K2 isoform present at high concentrations in the

> brain, kidney and pancreas, and is required for endothelial cell development. Mediates the conversion of phylloquinone (PK) into MK-4, probably by cleaving the side chain of phylloquinone (PK) to release 2-methyl-1,4-naphthoquinone (menadione; K3) and then

Source: prenylating it with geranylgeranyl pyrophosphate (GGPP) to form MK-4. Also plays a role in cardiovascular development independently of MK-4 biosynthesis, by acting as a Rabbit coenzyme Q10 biosyntetic enzyme: coenzyme Q10, also named ubiquinone, plays a

important antioxidant role in the cardiovascular system. Mediates biosynthesis of coenzyme Q10 in the Golgi membrane, leading to protect cardiovascular tissues from

NOS3/eNOS-dependent oxidative stress.

lgG

**Applications:** Gene ID:

**ELISA** UBIAD1

Uniprot **Recommended dilutions:** 

Q9Y5Z9

Synonyms:

UbiA prenyltransferase domain-containing protein 1 (EC 2.5.1) (Transitional epithelial

response protein 1), UBIAD1, TERE1

Immunogen:

Recombinant Human UbiA prenyltransferase domain-containing protein 1 protein (1-

82AA).

Storage:

Preservative: 0.03% Proclin 300. Constituents: 50% Glycerol, 0.01M PBS, pH 7.4

Product	<b>Images</b>
---------	---------------

N/A N/A