

DPY19L3 Antibody



PACO58212

Product Information

Size:

50ug

Reactivity:

Human

Source:

Rabbit

Isotype:

IgG

Applications:

ELISA, WB, IF

Recommended dilutions:

ELISA:1:2000-1:10000, WB:1:500-1:5000,
IF:1:200-1:500

Protein Background:

Probable C-mannosyltransferase that mediates C-mannosylation of tryptophan residues on target proteins.

Gene ID:

DPY19L3

Uniprot

Q6ZPD9

Synonyms:

Probable C-mannosyltransferase DPY19L3 (EC 2.4.1) (Dpy-19-like protein 3) (Protein dpy-19 homolog 3), DPY19L3


Immunogen:

Recombinant Human Probable C-mannosyltransferase DPY19L3 protein (587-697AA).


Storage:

Preservative: 0.03% Proclin 300. Constituents: 50% Glycerol, 0.01M PBS, pH 7.4

Product Images



Western Blot. Positive WB detected in: SH-SY5Y whole cell lysate, HepG2 whole cell lysate. All lanes: DPY19L3 antibody at 3.4 μ g/ml. Secondary. Goat polyclonal to rabbit IgG at 1/50000 dilution. Predicted band size: 84, 63 kDa. Observed band size: 84 kDa.



Immunofluorescence staining of HepG2 cells with PACO58212 at 1:200, counter-stained with DAPI. The cells were fixed in 4% formaldehyde, permeabilized using 0.2% Triton X-100 and blocked in 10% normal Goat Serum. The cells were then incubated with the antibody overnight at 4°C. The secondary antibody was Alexa Fluor 488-conjugated AffiniPure Goat Anti-Rabbit IgG(H+L).