AHNAK2 Antibody



PACO58272

Reactivity:

Product Information

Size: Protein Background:

50ug costamere, cytoplasmic vesicle membrane, cytosol, plasma membrane,

sarcolemma, T-tubule, Z disc.

Gene ID:

Human AHNAK2

Source: Uniprot

Rabbit Q8IVF2

Isotype:

lgG Synonyms:

Protein AHNAK2, AHNAK2, C14orf78 KIAA2019 **Applications:**

ELISA, IHC, IF


Recombinant Human Protein AHNAK2 protein (195-349AA).


ELISA:1:2000-1:10000, IHC:1:200-1:500,


Recommended dilutions:

IF:1:50-1:200 Preservative: 0.03% Proclin 300. Constituents: 50% Glycerol, 0.01M PBS, pH 7.4

Product Images







IHC image of PACO58272 diluted at 1:400 and staining in paraffinembedded human liver cancer performed on a Leica BondTM system. After dewaxing and hydration, antigen retrieval was mediated by high pressure in a citrate buffer (pH 6.0). Section was blocked with 10% normal goat serum 30min at RT. Then primary antibody (1% BSA) was incubated at 4°C overnight. The primary is detected by a biotinylated secondary antibody and visualized using an HRP conjugated SP system.

Immunofluorescence staining of A549 cells with PACO58272 at 1:133, counter-stained with DAPI. The cells were fixed in 4% formaldehyde, permeabilized using 0.2% Triton X-100 and blocked in 10% normal Goat Serum. The cells were then incubated with the antibody overnight at 4°C. The secondary antibody was Alexa Fluor 488-congugated AffiniPure Goat Anti-Rabbit IgG(H+L).

IHC image of PACO58272 diluted at 1:400 and staining in paraffinembedded human brain tissue performed on a Leica BondTM system. After dewaxing and hydration, antigen retrieval was mediated by high pressure in a citrate buffer (pH 6.0). Section was blocked with 10% normal goat serum 30min at RT. Then primary antibody (1% BSA) was incubated at 4°C overnight. The primary is detected by a biotinylated secondary antibody and visualized using an HRP conjugated SP system.