## **TFR2 Antibody**



## PACO58985

## **Product Information**

Size: Protein Background:

50ug Mediates cellular uptake of transferrin-bound iron in a non-iron dependent manner.

May be involved in iron metabolism, hepatocyte function and erythrocyte

Reactivity: differentiation.

Human Gene ID:

Source: TFR2

Rabbit Uniprot

**Isotype:** Q9UP52

lgG Synonyms:

**Applications:** Transferrin receptor protein 2 (TfR2), TFR2

ELISA, IHC, IF Immunogen:

**Recommended dilutions:** Recombinant Human Transferrin receptor protein 2 protein (534-605AA).


ELISA:1:2000-1:10000, IHC:1:200-1:500,


IF:1:50-1:200


Storage:

Preservative: 0.03% Proclin 300. Constituents: 50% Glycerol, 0.01M PBS, pH 7.4

## **Product Images**







IHC image of PACO58985 diluted at 1:200 and staining in paraffinembedded human liver cancer performed on a Leica BondTM system. After dewaxing and hydration, antigen retrieval was mediated by high pressure in a citrate buffer (pH 6.0). Section was blocked with 10% normal goat serum 30min at RT. Then primary antibody (1% BSA) was incubated at 4°C overnight. The primary is detected by a biotinylated secondary antibody and visualized using an HRP conjugated SP system.

Immunofluorescence staining of MCF-7 cells with PACO58985 at 1:66, counter-stained with DAPI. The cells were fixed in 4% formaldehyde, permeabilized using 0.2% Triton X-100 and blocked in 10% normal Goat Serum. The cells were then incubated with the antibody overnight at 4°C. The secondary antibody was Alexa Fluor 488-congugated AffiniPure Goat Anti-Rabbit IgG(H+L).

IHC image of PACO58985 diluted at 1:200 and staining in paraffinembedded human liver tissue performed on a Leica BondTM system. After dewaxing and hydration, antigen retrieval was mediated by high pressure in a citrate buffer (pH 6.0). Section was blocked with 10% normal goat serum 30min at RT. Then primary antibody (1% BSA) was incubated at 4°C overnight. The primary is detected by a biotinylated secondary antibody and visualized using an HRP conjugated SP system.