## **CSAD Antibody**



## PACO59704

**Product Information** 

Size: Protein Background:

50ug

Reactivity: Gene ID:

Human CSAD

Source: Uniprot

Rabbit Q9Y600

Isotype: Synonyms:

lgG Cysteine sulfinic acid, decarboxylase (EC 4.1.1.29) (Cysteine-sulfinate decarboxylase)

(Sulfinoalanine decarboxylase), CSAD, CSD

**Applications:** 

Immunogen:

ELISA, IHC

Recommended dilutions:


ELISA:1:2000-1:10000, IHC:1:500-1:1000


Storage:

Preservative: 0.03% Proclin 300. Constituents: 50% Glycerol, 0.01M PBS, pH 7.4

Recombinant Human Cysteine sulfinic acid, decarboxylase protein (344-493AA).

## **Product Images**





IHC image of PACO59704 diluted at 1:600 and staining in paraffinembedded human breast cancer performed on a Leica BondTM system. After dewaxing and hydration, antigen retrieval was mediated by high pressure in a citrate buffer (pH 6.0). Section was blocked with 10% normal goat serum 30min at RT. Then primary antibody (1% BSA) was incubated at 4°C overnight. The primary is detected by a biotinylated secondary antibody and visualized using an HRP conjugated SP system.

IHC image of PACO59704 diluted at 1:600 and staining in paraffinembedded human heart tissue performed on a Leica BondTM system. After dewaxing and hydration, antigen retrieval was mediated by high pressure in a citrate buffer (pH 6.0). Section was blocked with 10% normal goat serum 30min at RT. Then primary antibody (1% BSA) was incubated at 4°C overnight. The primary is detected by a biotinylated secondary antibody and visualized using an HRP conjugated SP system.