ZNF367 Antibody



PACO59944

Product Information

Recommended dilutions:

ELISA:1:2000-1:10000, WB:1:500-1:5000,

Size: Protein Background:

50ug Transcriptional activator. Isoform 1 may be involved in transcriptional activation of

erythroid genes.

Reactivity: Gene ID:

Human, Mouse ZNF367

Source:

Rabbit Uniprot

Q7RTV3 Isotype:

lgG Synonyms:


Zinc finger protein 367 (C2H2 zinc finger protein ZFF29), ZNF367, ZFF29 **Applications:**

ELISA, WB, IHC

Recombinant Human Zinc finger protein 367 protein (83-151AA).

Storage:

IHC:1:200-1:500 Preservative: 0.03% Proclin 300. Constituents: 50% Glycerol, 0.01M PBS, pH 7.4



Western Blot. Positive WB detected in: K562 whole cell lysate, NIH/3T3 whole cell lysate. All lanes: ZNF367 antibody at 6μ g/ml. Secondary. Goat polyclonal to rabbit IgG at 1/50000 dilution. Predicted band size: 39, 33 kDa. Observed band size: 39 kDa.

IHC image of PACO59944 diluted at 1:400 and staining in paraffinembedded human glioma performed on a Leica BondTM system. After dewaxing and hydration, antigen retrieval was mediated by high pressure in a citrate buffer (pH 6.0). Section was blocked with 10% normal goat serum 30min at RT. Then primary antibody (1% BSA) was incubated at 4°C overnight. The primary is detected by a biotinylated secondary antibody and visualized using an HRP conjugated SP system.