
PACO59948



Product Information	
Size:	Protein Background:
50ug	Plays a role as a receptor for the recognition of MHC class I HLA-E molecules by NK
Reactivity:	cells and some cytotoxic T-cells.
Human	Gene ID:
Source:	KLRC3
Rabbit	Uniprot
	Q07444
Isotype:	Synonyms:
lgG	NKG2-E type II integral membrane protein (NK cell receptor E) (NKG2-E-activating NK
Applications:	receptor), KLRC3, NKG2E
ELISA, IHC, IF	Immunogen:
Recommended dilutions:	Recombinant Human NKG2-E type II integral membrane protein (94-240AA).
ELISA:1:2000-1:10000, IHC:1:200-1:500, IF:1:50-1:200	Storage:
11.1.50 1.200	Preservative: 0.03% Proclin 300. Constituents: 50% Glycerol, 0.01M PBS, pH 7.4



IHC image of PACO59948 diluted at 1:400 and staining in paraffinembedded human tonsil tissue performed on a Leica BondTM system. After dewaxing and hydration, antigen retrieval was mediated by high pressure in a citrate buffer (pH 6.0). Section was blocked with 10% normal goat serum 30min at RT. Then primary antibody (1% BSA) was incubated at 4°C overnight. The primary is detected by a biotinylated secondary antibody and visualized using an HRP conjugated SP system.

Immunofluorescence staining of Hela cells with PACO59948 at 1:133, counter-stained with DAPI. The cells were fixed in 4% formaldehyde, permeabilized using 0.2% Triton X-100 and blocked in 10% normal Goat Serum. The cells were then incubated with the antibody overnight at 4°C. The secondary antibody was Alexa Fluor 488-congugated AffiniPure Goat Anti-Rabbit IgG(H+L).