

# DAZAP2 Antibody



PACO60028

---

## Product Information

**Size:**

50ug

**Reactivity:**

Human, Rat

**Source:**

Rabbit

**Isotype:**

IgG

**Applications:**

ELISA, WB, IHC, IF

**Recommended dilutions:**

ELISA:1:2000-1:10000, WB:1:500-1:5000,  
IHC:1:200-1:500, IF:1:50-1:200

**Protein Background:**

nuclear speck, identical protein binding, WW domain binding.

**Gene ID:**

DAZAP2

**Uniprot**

Q15038

**Synonyms:**

DAZ-associated protein 2 (Deleted in azoospermia-associated protein 2), DAZAP2, KIAA0058

**Immunogen:**




Recombinant Human DAZ-associated protein 2 protein (1-168AA).

**Storage:**

Preservative: 0.03% Proclin 300. Constituents: 50% Glycerol, 0.01M PBS, pH 7.4

## Product Images

---



Western Blot. Positive WB detected in: HeLa whole cell lysate, A375 whole cell lysate, MCF-7 whole cell lysate, A549 whole cell lysate, SH-SY5Y whole cell lysate, Rat brain tissue. All lanes: DAZAP2 antibody at 3.5 $\mu$ g/ml. Secondary. Goat polyclonal to rabbit IgG at 1/50000 dilution. Predicted band size: 18, 17, 16, 14, 10, 23 kDa. Observed band size: 18 kDa.

Immunofluorescence staining of MCF-7 cells with PACO60028 at 1:133, counter-stained with DAPI. The cells were fixed in 4% formaldehyde, permeabilized using 0.2% Triton X-100 and blocked in 10% normal Goat Serum. The cells were then incubated with the antibody overnight at 4°C. The secondary antibody was Alexa Fluor 488-conjugated AffiniPure Goat Anti-Rabbit IgG(H+L).

IHC image of PACO60028 diluted at 1:400 and staining in paraffin-embedded human prostate tissue performed on a Leica Bond™ system. After dewaxing and hydration, antigen retrieval was mediated by high pressure in a citrate buffer (pH 6.0). Section was blocked with 10% normal goat serum 30min at RT. Then primary antibody (1% BSA) was incubated at 4°C overnight. The primary is detected by a biotinylated secondary antibody and visualized using an HRP conjugated SP system.