
RNF220 Antibody

PACO60088



| Product Information | |
|--|--|
| Size: | Protein Background: |
| 50ug | E3 ubiquitin-protein ligase that promotes the ubiquitination and proteasomal |
| Reactivity: | degradation of SIN3B. Independently of its E3 ligase activity, acts as a CTNNB1 stabilizer through USP7-mediated deubiquitination of CTNNB1 promoting Wnt signaling. |
| Human | Gene ID: |
| Source: | RNF220 |
| Rabbit | Uniprot |
| lsotype: | Q5VTB9 |
| lgG | Synonyms: |
| Applications: | E3 ubiquitin-protein ligase RNF220 (EC 2.3.2.27) (RING finger protein 220) (RING-type |
| ELISA, IHC | E3 ubiquitin transferase RNF220), RNF220, C1orf164 |
| Recommended dilutions: | Immunogen: |
| ELISA:1:2000-1:10000, IHC:1:500-1:1000 | Recombinant Human E3 ubiquitin-protein ligase RNF220 protein (384-472AA). |
| | Storage: |
| | Preservative: 0.03% Proclin 300. Constituents: 50% Glycerol, 0.01M PBS, pH 7.4 |



IHC image of PACO60088 diluted at 1:500 and staining in paraffinembedded human prostate cancer performed on a Leica BondTM system. After dewaxing and hydration, antigen retrieval was mediated by high pressure in a citrate buffer (pH 6.0). Section was blocked with 10% normal goat serum 30min at RT. Then primary antibody (1% BSA) was incubated at 4°C overnight. The primary is detected by a biotinylated secondary antibody and visualized using an HRP conjugated SP system.