

PACO61814

---

## Product Information

**Size:**

50ul

**Reactivity:**

Mycobacterium kansasii

**Source:**

Rabbit

**Isotype:**

IgG

**Applications:**

ELISA, WB

**Recommended dilutions:**

ELISA:1:2000-1:10000, WB:1:500-1:5000

**Protein Background:**

The antigen 85 proteins (FbpA, FbpB, FbpC) are responsible for the high affinity of mycobacteria for fibronectin, a large adhesive glycoprotein, which facilitates the attachment of *M. tuberculosis* to murine alveolar macrophages (AMs). They also help to maintain the integrity of the cell wall by catalyzing the transfer of mycolic acid, to cell wall arabinogalactan and through the synthesis of alpha, alpha-trehalose dimycolate (TDM, cord factor). They catalyze the transfer of a mycoloyl residue from one molecule of alpha, alpha-trehalose monomycolate (TMM) to another TMM, leading to the formation of TDM.

**Gene ID:**

fbpB

**Uniprot**

P21160

**Synonyms:**

Diacylglycerol acyltransferase/mycolyltransferase Ag85B (DGAT) (EC 2.3.1.122) (EC 2.3.1.20) (30 kDa extracellular protein) (Acyl-CoA: diacylglycerol acyltransferase) (Antigen 85 complex B) (85B) (Ag85B) (Extracellular alpha-antigen) (Fibronectin-binding protein B) (Fbps B), fbpB

**Immunogen:**


Recombinant *Mycobacterium kansasii* Diacylglycerol acyltransferase/mycolyltransferase Ag85B protein (41-325AA).

**Storage:**

Preservative: 0.03% Proclin 300. Constituents: 50% Glycerol, 0.01M PBS, pH 7.4

## Product Images

---



Western Blot. Positive WB detected in Recombinant protein. All lanes: fbpB antibody at 3.2 $\mu$ g/ml. Secondary. Goat polyclonal to rabbit IgG at 1/50000 dilution. Predicted band size: 50 kDa. Observed band size: 50 kDa.