

Product Information

Size:

50ul

Reactivity:

Human

Source:

Rabbit

Isotype:

IgG

Applications:

ELISA, WB

Recommended dilutions:

ELISA:1:2000-1:10000, WB:1:1000-1:5000

Protein Background:

Deubiquitinating enzyme that removes conjugated ubiquitin from specific proteins to regulate different cellular processes that may include cell proliferation, progression through the cell cycle, apoptosis, cell migration, and the cellular response to viral infection.

Gene ID:

USP17L1

Uniprot

Q7RTZ2

Synonyms:

Ubiquitin carboxyl-terminal hydrolase 17-like protein 1 (EC 3.4.19.12) (Deubiquitinating enzyme 17-like protein 1) (Ubiquitin thioesterase 17-like protein 1) (Ubiquitin-specific-processing protease 17-like protein 1), USP17L1, USP17L USP17L1P


Immunogen:

Recombinant Human Ubiquitin carboxyl-terminal hydrolase 17-like protein 1 protein (19-235AA).


Storage:

Preservative: 0.03% Proclin 300. Constituents: 50% Glycerol, 0.01M PBS, pH 7.4

Product Images



Western Blot. Positive WB detected in: U87 whole cell lysate. All lanes: USP17L1 antibody at 1:2000. Secondary. Goat polyclonal to rabbit IgG at 1/50000 dilution. Predicted band size: 60 kDa. Observed band size: 60 kDa.



Western Blot. Positive WB detected in: HeLa whole cell lysate, U87 whole cell lysate. All lanes: USP17L1 antibody at 1:2000. Secondary. Goat polyclonal to rabbit IgG at 1/50000 dilution. Predicted band size: 60 kDa. Observed band size: 60 kDa.