## APOC2 Antibody, FITC conjugated

## PACO63941



| Product Information    |  |
|------------------------|--|
| Size:                  | Protein Background:  |
| 50ul                   | Component of chylomicrons, very low-density lipoproteins (VLDL), low-density   |
| Reactivity:            | lipoproteins (LDL), and high-density lipoproteins (HDL) in plasma. Plays an important role in lipoprotein metabolism as an activator of lipoprotein lipase. Both |
| Human                  | proapolipoprotein C-II and apolipoprotein C-II can activate lipoprotein lipase. In normolipidemic individuals, it is mainly distributed in the HDL, whereas in   |
| Source:                | hypertriglyceridemic individuals, predominantly found in the VLDL and LDL.   |
| Rabbit                 | Gene ID:   |
| lsotype:               | APOC2  |
| IgG                    | Uniprot  |
| Applications:          | P02655   |
| ELISA                  | Synonyms:  |
| Recommended dilutions: | Apolipoprotein C-II (Apo-CII) (ApoC-II) (Apolipoprotein C2) [Cleaved into:<br>Proapolipoprotein C-II (ProapoC-II)], APOC2, APC2                                  |
|                        | Immunogen:   |
|                        | Peptide sequence from Human Apolipoprotein C-II protein (20-38AA).   |
|                        | Storage:   |
|                        | Preservative: 0.03% Proclin 300. Constituents: 50% Glycerol, 0.01M PBS, pH 7.4   |

N/A N/A