

## **Technical Manual**

# **ChromaDazzle alpha-L-Fucosidase Activity Assay Kit**

**Catalogue Code: BA0023**

**Pack Size: 100 assays**

**Research Use Only**



**Standard Preparation:** Mix 10  $\mu\text{L}$  of 12.5 mM nitrophenol standard with 490  $\mu\text{L}$  dH<sub>2</sub>O to make 250  $\mu\text{M}$  standard.

No	250 $\mu\text{M}$ STD + dH <sub>2</sub> O	Vol ( $\mu\text{L}$ )	Nitrophenol ( $\mu\text{M}$ )
1	200 $\mu\text{L}$ + 0 $\mu\text{L}$	200	250
2	120 $\mu\text{L}$ + 80 $\mu\text{L}$	200	150
3	60 $\mu\text{L}$ + 140 $\mu\text{L}$	200	75
4	0 $\mu\text{L}$ + 200 $\mu\text{L}$	200	0

### Reaction Preparation:

1. Transfer 100  $\mu\text{L}$  of each standard (OD<sub>STD</sub>) into wells of a clear flat bottom 96-well plate.
2. Transfer 20  $\mu\text{L}$  of each sample into separate wells. Add 80  $\mu\text{L}$  Substrate to each sample well. Tap plate briefly to mix.
3. Incubate at 25°C or desired temperature for 20 minutes. Add 100  $\mu\text{L}$  of Stop Reagent to each well. Tap plate briefly to mix.
4. Read OD<sub>405nm</sub>.

*Note: If your sample is colored or opaque, then a sample blank (OD<sub>BLANK</sub>) will be needed. Add 20  $\mu\text{L}$  of sample to a well, and add 80  $\mu\text{L}$  of dH<sub>2</sub>O. After incubation add 100  $\mu\text{L}$  Stop Reagent.*

### CALCULATION


Subtract blank OD (water, #4) from the standard OD values and plot the  $\Delta\text{OD}$  against standard concentrations. Determine the Slope and use the following equation to calculate alpha-Fucosidase activity:

$$\begin{aligned} \text{AFU Activity} &= \frac{\text{OD}_{\text{SAMPLE}} - \text{OD}_{\text{BLANK}}}{\text{Time} \cdot \text{Slope}} \times \frac{\text{Reaction Vol } (\mu\text{L})}{\text{Sample Vol } (\mu\text{L})} \times n \\ &= \frac{\text{OD}_{\text{SAMPLE}} - \text{OD}_{\text{BLANK}}}{\text{Slope}} \times \frac{1}{4} \times n \quad (\text{U/L}) \end{aligned}$$


where OD<sub>SAMPLE</sub> is the OD<sub>405nm</sub> value for each sample and OD<sub>BLANK</sub> is the OD<sub>405nm</sub> value of the water (standard #4) or the sample blank if one was used. Slope is the slope of the linear regression fit of the standard points and Time is the reaction time (20 min). Reaction Vol and Sample Vol are 100  $\mu\text{L}$  and 20  $\mu\text{L}$ , respectively.  $n$  is the dilution factor.

*Unit definition:* 1 Unit (U) of AFU will catalyze the conversion of 1  $\mu\text{mole}$  of 4-Nitrophenyl-alpha-L-fucopyranoside to 4-Nitrophenol and alpha-L-Fucose per min at 25°C and pH 5.3.

*Note: If sample AFU activity exceeds 100 U/L, either use a shorter reaction time or dilute samples in water and repeat the assay. For samples with AFU activity < 5 U/L, the incubation time can be extended up to 40 minutes for greater sensitivity.*



**Titration Curve**



**Standard Curve**

#### **MATERIALS REQUIRED, BUT NOT PROVIDED**

Pipetting devices and accessories (e.g. multi-channel pipettor), clear flat-bottom 96-well plates (e.g. VWR cat# 82050-760), centrifuge tubes and plate reader.

#### **LITERATURE**

1. GIARDINA M.G. ET AL. (1992) SERUM ALPHA-L-FUCOSIDASE. A USEFUL MARKER IN THE DIAGNOSIS OF HEPATOCELLULAR CARCINOMA. *CANCER*. 70:1044-1048
2. Fernandez-Rodriguez AD. Et al. (2000) Value of the serum alpha-L-fucosidase activity in the diagnosis of colorectal cancer. *Oncology* 59(4): 310-316.
3. Alhadef JA et al. (1975) Human liver alpha-L-fucosidase. Purification, characterization, and immunochemical studies. *Journal Biol. Chem.* 250: 7106-7113.

### **Contact Details**

**Dublin, Ireland**

**Email:** [info@assaygenie.com](mailto:info@assaygenie.com)

**Web:** [www.assaygenie.com](http://www.assaygenie.com)

**Technical Support:** [Techsupport@assaygenie.com](mailto:Techsupport@assaygenie.com)