

# Recombinant Protein Technical Manual Recombinant Human IL3RA/CD123 Protein (Fc Tag)

**RPES1474** 

#### **Product Data:**

**Product SKU:** RPES1474 **Size:** 10μg

Species: Human Cells

**Uniprot:** P26951

### **Protein Information:**

Molecular Mass: 59.9 kDa

AP Molecular Mass: 8000 kDa

Tag: C-Fc

**Bio-activity:** 

**Purity:** > 95% as determined by reducing SDS-PAGE.

**Endotoxin:**  $< 1.0 \text{ EU per } \mu\text{g}$  as determined by the LAL method.

**Storage:** Lyophilized protein should be stored at < -20°C, though stable at room

temperature for 3 weeks. Reconstituted protein solution can be stored at 4-7°C for 2-7 days. Aliquots of reconstituted samples are stable at < -20°C for 3 months.

**Shipping:** This product is provided as lyophilized powder which is shipped with ice packs.

**Formulation:** Lyophilized from a 0.2 μm filtered solution of PBS, pH7.4.

**Reconstitution:** Please refer to the printed manual for detailed information.

**Application:** 

Synonyms: Interleukin-3 receptor subunit alpha; IL-3 receptor subunit alpha; IL-3R subunit

alpha; IL-3R-alpha; IL-3RA

## Immunogen Information:

Sequence: Thr19-Arg305

## **Background**:

CD123, also known as Interleukin-3 receptor subunit alpha, belongs to the type I cytokine receptor family. In mouse, there are two classes of high-affinity IL3 receptors. One contains an IL3-specific beta subunit and the other contains the beta subunit also shared by high-affinity IL5 and GM-CSF receptors. CD123 stimulates the proliferation and differentiation of hemopoietic cells including the pluripotent hematopoietic stem cells as well as various lineage-committed cells. CD123 is a heterodimer consisting of an alpha and a beta subunit. The alpha subunit alone binds IL-3 with low affinity. The beta subunit does not bind IL-3 by itself but is required for the high-affinity binding of IL-3 to the heterodimeric receptor complex.