



# Recombinant Protein Technical Manual

**Recombinant Mouse AKT3 Protein (aa 106-479)(Active)**  
RPES1835

## Product Data:

**Product SKU:** RPES1835

**Size:** 20µg

**Species:** Mouse

**Expression host:** Baculovirus-Insect Cells

**Uniprot:** Q9WUA6

## Protein Information:

**Molecular Mass:** 43.4 kDa

**AP Molecular Mass:** 46 kDa

### Tag:

**Bio-activity:** The specific activity was determined to be 20 nmol/min/mg using synthetic GSK3-derived peptide (R11-SGRARTSSFAEPGGK) as substrate.

**Purity:** > 95 % as determined by SDS-PAGE

**Endotoxin:** < 1.0 EU per µg of the protein as determined by the LAL method.

**Storage:** Store at < -20°C, stable for 6 months. Please minimize freeze-thaw cycles.

**Shipping:** This product is provided as liquid. It is shipped at frozen temperature with blue ice/gel packs. Upon receipt, store it immediately at < -20°C.

**Formulation:** Supplied as sterile 20mM Tris, 500mM NaCl, 10% glycerol, pH 7.4

**Reconstitution:** Please refer to the printed manual for detailed information.

### Application:

**Synonyms:** AI851531;D930002M15Rik;Nmf350

## Immunogen Information:

**Sequence:** Ala106-Glu479

## Background:

v-akt murine thymoma viral oncogene homolog 3 (AKT3), also known as PKB-GAMMA, with AKT1/PKBalpha, AKT2/PKBbeta, are the members of Akt kinase family, share extensive structural similarity and perform common as well as unique functions within cells. The Akt signaling cascade initiates at the cell surface when growth factors or other extracellular stimuli activate phosphoinositide 3-kinase (PI3K). AKT3 was discovered to be the predominant isoform activated in sporadic melanomas. Levels of activity increased during melanoma progression with metastatic melanomas having the highest activity. Although mechanisms of AKT3 activation remain to be fully characterized, overexpression of AKT3 and decreased PTEN activity play important roles in this process. Targeted reduction of AKT3 activity decreased survival of melanoma tumor cells leading to inhibition of tumor development, which may be therapeutically effective for shrinking tumors in melanoma patients. AKT2 and AKT3 play an important role in the viability of human malignant glioma cells. Targeting AKT2 and AKT3 may hold promise for the treatment of patients with gliomas.