

Recombinant Protein Technical Manual

Recombinant Human Pro-Neuregulin/NRG1--β1 Protein (aa 1-246, His Tag)(Active) RPES2788

Product Data:

Product SKU: RPES2788 **Size:** 10μg

Species: Human Expression host: E. coli

Uniprot: Q02297-6

Protein Information:

Molecular Mass: 29.0 kDa

AP Molecular Mass: 38 kDa

Tag: N-6His

Bio-activity: Immobilized Human NRG1Beta-His at 10μg/ml(100 μl/well) can bind Human

HER3-Fc(Cat: PKSH033438). The ED50 of Human Glypican-3-His is 5.18 ug/ml.

Purity: > 95 % as determined by reducing SDS-PAGE.

Endotoxin: < 1.0 EU per μg as determined by the LAL method.

Storage: Lyophilized proteins are stable for up to 12 months when stored at -20 to -80°C.

Reconstituted protein solution can be stored at 4-8°C for 2-7 days. Aliquots of

reconstituted samples are stable at < -20°C for 3 months.

Shipping: This product is provided as lyophilized powder which is shipped with ice packs.

Formulation: Lyophilized from a 0.2 μm filtered solution of 4mM HCl.

Reconstitution: Please refer to the printed manual for detailed information.

Application: Functional ELISA

Synonyms: Pro-neuregulin; Neuregulin beta 1; NRG1-beta 1; HRG1-beta 1; EGF; NRG1; GGF;

HGL; HRGA; NDF; SMDF;

Immunogen Information:

Sequence: Met 1-Lys246

Background:

neuregulin (heregulin, NRG1) is a member of neuregulin family, which is comprised of four genes that encode a large number of secreted or membrane-bound isoforms. All family members share an EGF-like domain that interacts with the ErbB family of tyrosine kinase receptors. NRG1 isoforms can be classified into type I, type II and type III isoforms. NRG1 directs ligand for ERBB3 and ERBB4 tyrosine kinase receptors, concomitantly recruits ERBB1 and ERBB2 coreceptors, resulting in ligand-stimulated tyrosine phosphorylation and activation of the ERBB receptors. NRG proteins show distinct spatial and temporal expression patterns and play important roles during development of both the nervous system and the heart.