

Recombinant Protein Technical Manual

Recombinant Human Carbonic Anhydrase 4/CA4 Protein (His Tag)(Active) **RPES2882**

Species: Human

Size: 10µg

Expression host: HEK293 Cells

Uniprot: NP_000708.1

Molecular Mass:	31.7 kDa
AP Molecular Mass:	30 kDa
Tag:	C-His
Bio-activity:	Measured by its esterase activity. The specific activity is >2 pmoles/min/ μ g.
Purity:	> 96 % as determined by reducing SDS-PAGE.
Endotoxin:	< 1.0 EU per μg as determined by the LAL method.
Storage:	Lyophilized proteins are stable for up to 12 months when stored at -20 to -80°C. Reconstituted protein solution can be stored at 4-8°C for 2-7 days. Aliquots of reconstituted samples are stable at < -20°C for 3 months.
Shipping:	This product is provided as lyophilized powder which is shipped with ice packs.
Formulation:	Lyophilized from sterile PBS, pH 7.4
Reconstitution:	Please refer to the printed manual for detailed information.
Application:	
Synonyms:	Carbonic Anhydrase 4; Carbonate Dehydratase IV; Carbonic Anhydrase IV; CA-IV; CA4;CAIV;Car4;RP17

Sequence: Met 1-Lys283

Background:

The carbonic anhydrases (or carbonate dehydratases) are classified as metalloenzyme for its zinc ion prosthetic group and form a family of enzymes that catalyze the rapid interconversion of carbon dioxide and water to bicarbonate and protons, a reversible reaction that takes part in maintaining acid-base balance in blood and other tissues. The carbonic anhydrasekl (CA) family consists of at least 11 enzymatically active members and a few inactive homologous proteins. Carbonic anhydrase IV (CAIV) is a membrane-associated enzyme anchored to plasma membrane surfaces by a phosphatidylinositol glycan linkage. CAIV is a high-activity isozyme in CO2 hydration comparable to that of CAII. Furthermore, CAIV is more active in HCO3-dehydration than is CAII. However, the esterase activity of CAIV is decreased 150-fold compared to CAII.