



Recombinant Protein Technical Manual
Recombinant Human G-CSF/CSF3 Protein (isoform
b)(Active)
RPES2904

Product Data:

Product SKU: RPES2904

Size: 5µg

Species: Human

Expression host: E. coli

Uniprot: NP_757373.1

Protein Information:

Molecular Mass: 18.7 kDa

AP Molecular Mass: 18 kDa

Tag:

Bio-activity: Measured in a cell proliferation assay using NFS-60 mouse myelogenous leukemia lymphoblast cells. The ED50 for this effect is typically 0.04-0.2 ng/mL.

Purity: > 90 % as determined by reducing SDS-PAGE.

Endotoxin: Please contact us for more information.

Storage: Lyophilized proteins are stable for up to 12 months when stored at -20 to -80°C. Reconstituted protein solution can be stored at 4-8°C for 2-7 days. Aliquots of reconstituted samples are stable at < -20°C for 3 months.

Shipping: This product is provided as lyophilized powder which is shipped with ice packs.

Formulation: Lyophilized from sterile sterile PBS, pH 7.4

Reconstitution: Please refer to the printed manual for detailed information.

Application: Cell Culture

Synonyms: Granulocyte Colony-Stimulating Factor; G-CSF; Pluripoietin; Filgrastim; Lenograstim; CSF3; C17orf33; GCSF

Immunogen Information:

Sequence: Ala30-Pro204

Background:

Granulocyte-colony stimulating factor (G-CSF) is a growth factor and an essential cytokine belonging to the CSF family of hormone-like glycoproteins. It is produced by numerous cell types including immune and endothelial cells. G-CSF binding to its receptor G-CSF-R which belongs to the cytokine receptor type I family depends on the interaction of alpha-helical motifs of the former and two fibronectin type III as well as an immunoglobulin-like domain of the latter. Recent animal studies have also revealed that G-CSF activates multiple signaling pathways, such as Akt and also the Janus family kinase-2 and signal transducer and activation of transcription-3 (Jak2-STAT3) pathway, thereby promoting survival, proliferation, differentiation and mobilisation of haematopoietic stem and progenitor cells. G-CSF is a cytokine that have been demonstrated to improve cardiac function and perfusion in myocardial infarction. And it was initially evaluated as a stem cell mobilizer and erythropoietin as a cytoprotective agent. G-CSF prevents left ventricular remodeling after myocardial infarction by decreasing cardiomyocyte death and by increasing the number of blood vessels, suggesting the importance of direct actions of G-CSF on the myocardium rather than through mobilization and differentiation of stem cells. Accordingly, recombinant human (rh)G-CSF has been extensively used in clinical haematology and oncology to enable bone marrow transplantation or to treat chemotherapy-associated neutropenia. In preclinical study, G-CSF improved cardiac function and perfusion by angiomyogenesis and protection of cardiomyocytes in myocardial infarction.