

Product Information	Immunogen information
Product SKU:	Background:
CAB22295	Histones are basic nuclear proteins that are responsible for the nucleosome structure of the
Gene ID:	chromosomal fiber in eukaryotes. Nucleosomes consist of approximately 146 bp of DNA wrapped around a histone octamer composed of pairs of each of the four core histones (H2A,
82908350	H2B, H3, and H4). The chromatin fiber is further compacted through the interaction of a linker histone, H1, with the DNA between the nucleosomes to form higher order chromatin structures.
Observed MW:	This gene is intronless and encodes a replication-dependent histone that is a member of the histone H3 family. Transcripts from this gene lack polyA tails; instead, they contain a palindromic
17kDa	termination element. This gene is located separately from the other H3 genes that are in the histone gene cluster on chromosome 6p22-p21.3.
Calculated MW:	historie gene cluster on chromosome op22-p21.3.
15kDa	Size:
Category:	20uL, 100uL
Primary Antibody	Applications:
Uniprot:	Western blotting Immunofluorescence
-	Synonyms:
Q16695P68431	H3.4; H3/g; H3FT; H3t; HIST3H3; Histone H3; HIST1H3A; TriMethyl-Histone H3-K9
Reactivity:	Immunogen:
Human, Mouse, Rat, Other (Wide Range Predicted)	A synthetic trimethylated peptide around K9 of human Histone H3 (NP_003520.1).
Purification Method:	Recommended Dilution:
Affinity purification	WB 1:2000 - 1:20000 IF/ICC 1:50 - 1:200 ChIP 5µg antibody for 5µg-10µg of ChromatinChIP-seq 1:20 - 1:100 DB 1:2000 - 1:20000 CUT&Tag 10 <sup>5</sup> cells /1 µg
Source:	Storage:
Rabbit	Store at -20°C. Avoid freeze / thaw cycles.Buffer: PBS with 0.05% proclin300, 0.05% BSA, 50% glycerol, pH7.3.
lsotype:	
lgG	