## Nanodisc Human OR8B8-Strep Protein



## HDFP1146

## **Product Information**

Product SKU:	HDFP1146	Expression Host:	HEK293		Size:	10µg
Target:	OR8B8	Tag:	C-Flag&St	rep Tag		
Additional Infor	mation					
<b>Conjugate</b> :	Unconjugat	ed <b>Unip</b>	orot ID:	Q15620		
Molecular Wei	<b>ght:</b> The human	full length OR8B8-Sti	rep protein l	has a MW o	of 34.5 kDa	

## **Protein Information**

Background:	Olfactory receptors interact with odorant molecules in the nose, to initiate a neuronal
	response that triggers the perception of a smell. The olfactory receptor proteins are
	members of a large family of G-protein-coupled receptors (GPCR) arising from single
	coding-exon genes. Olfactory receptors share a 7-transmembrane domain structure
	with many neurotransmitter and hormone receptors and are responsible for the
	recognition and G protein-mediated transduction of odorant signals. The olfactory
	receptor gene family is the largest in the genome. The nomenclature assigned to the
	olfactory receptor genes and proteins for this organism is independent of other
	organisms. [provided by RefSeq, Jul 2008]
Synonyms:	TPCR85
Protein Description:	Human OR8B8-Strep full length protein-synthetic nanodisc
Protein Description: Formulation:	Human OR8B8-Strep full length protein-synthetic nanodisc Lyophilized from nanodisc solubilization buffer (20 mM Tris-HCl, 150 mM NaCl, pH
•	
•	Lyophilized from nanodisc solubilization buffer (20 mM Tris-HCl, 150 mM NaCl, pH
•	Lyophilized from nanodisc solubilization buffer (20 mM Tris-HCl, 150 mM NaCl, pH 8.0). Normally 5% – 8% trehalose is added as protectants before lyophilization. Please
•	Lyophilized from nanodisc solubilization buffer (20 mM Tris-HCl, 150 mM NaCl, pH 8.0). Normally 5% – 8% trehalose is added as protectants before lyophilization. Please see Certificate of Analysis for specific instructions. Do not use solvents with a pH
•	Lyophilized from nanodisc solubilization buffer (20 mM Tris-HCl, 150 mM NaCl, pH 8.0). Normally 5% – 8% trehalose is added as protectants before lyophilization. Please see Certificate of Analysis for specific instructions. Do not use solvents with a pH below 6.5 or those containing high concentrations of divalent metal ions (greater
Formulation:	Lyophilized from nanodisc solubilization buffer (20 mM Tris-HCl, 150 mM NaCl, pH 8.0). Normally 5% – 8% trehalose is added as protectants before lyophilization. Please see Certificate of Analysis for specific instructions. Do not use solvents with a pH below 6.5 or those containing high concentrations of divalent metal ions (greater than 5 mM) in subsequent experiments.

**Storage & Shipping**: Store at -20°C to -80°C for 12 months in lyophilized form. After reconstitution, if not intended for use within a month, aliquot and store at -80°C (Avoid repeated freezing and thawing). Lyophilized proteins are shipped at ambient temperature.