## Nanodisc Human S1PR5-Strep Protein



## HDFP1187

## **Product Information**

<b>Product SKU</b> :	HDF	P1187	Expression Ho	<b>st</b> : HEK293		Size:	10µg
Target:	S1PR5		Tag:	C-Flag&S	Strep Tag		
Additional Infor	matio	'n					
Conjugate:		Unconjugate	ed <b>U</b>	niprot ID:	Q9H228		
Molecular Wei	ght:	The human f	ull length S1PR5	-Strep protein	has a MW c	f 41.8 kDa	
Protein Informa	tion						
Background <sup>.</sup>		The lyces	phingolinid sphi	ingocino 1-nh	ocobato (S <sup>r</sup>	1P) regulator	coll proliforation

Background	The lysosphingolipid sphingosine 1-phosphate (S1P) regulates cell proliferation,
	apoptosis, motility, and neurite retraction. Its actions may be both intracellular as a
	second messenger and extracellular as a receptor ligand. S1P and the structurally
	related lysolipid mediator lysophosphatidic acid (LPA) signal cells through a set of G
	protein-coupled receptors known as EDG receptors. Some EDG receptors (e.g., EDG1;
	MIM 601974) are S1P receptors; others (e.g., EDG2; MIM 602282) are LPA
	receptors.[supplied by OMIM, Mar 2008]
Synonyms:	EDG8, Edg-8, S1P5, SPPR-1, SPPR-2
Protein Description:	Human S1PR5-Strep full length protein-synthetic nanodisc
<b>Formulation</b> :	Lyophilized from nanodisc solubilization buffer (20 mM Tris-HCl, 150 mM NaCl, pH
Formulation:	Lyophilized from nanodisc solubilization buffer (20 mM Tris-HCl, 150 mM NaCl, pH 8.0). Normally 5% – 8% trehalose is added as protectants before lyophilization. Please
Formulation:	
Formulation:	8.0). Normally 5% – 8% trehalose is added as protectants before lyophilization. Please
Formulation:	8.0). Normally 5% – 8% trehalose is added as protectants before lyophilization. Please see Certificate of Analysis for specific instructions. Do not use solvents with a pH
Formulation: Protein Pathways:	8.0). Normally 5% – 8% trehalose is added as protectants before lyophilization. Please see Certificate of Analysis for specific instructions. Do not use solvents with a pH below 6.5 or those containing high concentrations of divalent metal ions (greater
	8.0). Normally 5% – 8% trehalose is added as protectants before lyophilization. Please see Certificate of Analysis for specific instructions. Do not use solvents with a pH below 6.5 or those containing high concentrations of divalent metal ions (greater than 5 mM) in subsequent experiments.

**Storage & Shipping**: Store at -20°C to -80°C for 12 months in lyophilized form. After reconstitution, if not intended for use within a month, aliquot and store at -80°C (Avoid repeated freezing and thawing). Lyophilized proteins are shipped at ambient temperature.