

## **HDFP1262**

## **Product Information**

Product SKU:	HDFP1262	Expression Host:	HEK293		Size:	10µg
Target:	PLM	Tag:	C-Flag&St	rep Tag		
Additional Infor	mation					
<b>Conjugate</b> :	Unconjugat	ted Unip	prot ID:	O00168		
Molecular Wei	ght: The human	The human full length PLM-Strep protein has a MW of 10.4 kDa				

## **Protein Information**

Background:	: This gene encodes a member of a family of small membrane proteins that share a				
	35-amino acid signature sequence domain, beginning with the sequence PFXYD and				
	containing 7 invariant and 6 highly conserved amino acids. The approved hum				
	gene nomenclature for the family is FXYD-domain containing ion transport regulator				
	Mouse FXYD5 has been termed RIC (Related to Ion Channel). FXYD2, also known as				
	the gamma subunit of the Na,K-ATPase, regulates the properties of that enzyme.				
	FXYD1 (phospholemman), FXYD2 (gamma), FXYD3 (MAT-8), FXYD4 (CHIF), and FXYD5				
	(RIC) have been shown to induce channel activity in experimental expression systems.				
	Transmembrane topology has been established for two family members (FXYD1 and				
	FXYD2), with the N-terminus extracellular and the C-terminus on the cytoplasmic side				
	of the membrane. The protein encoded by this gene is a plasma membrane substrate				
	for several kinases, including protein kinase A, protein kinase C, NIMA kinase, and				
	myotonic dystrophy kinase. It is thought to form an ion channel or regulate ion				
	channel activity. Transcript variants with different 5' UTR sequences have been				
	described in the literature. [provided by RefSeq, Jul 2008]				
Synonyms:	PLM				
Protein Description:	Human PLM-Strep full length protein-synthetic nanodisc				
<b>Formulation</b> :	Lyophilized from nanodisc solubilization buffer (20 mM Tris-HCl, 150 mM NaCl, pH				
	8.0). Normally 5% – 8% trehalose is added as protectants before lyophilization. Please				

	see Certificate of Analysis for specific instructions. Do not use solvents with a pH		
	below 6.5 or those containing high concentrations of divalent metal ions (greater		
	than 5 mM) in subsequent experiments.		
Protein Pathways:	-		
Protein Families:	Ion Channels: Other.		
Usage:	Research use only		
Storage & Shipping:	Store at -20°C to -80°C for 12 months in lyophilized form. After reconstitution, if not		
	intended for use within a month, aliquot and store at -80°C (Avoid repeated freezing		
	and thawing). Lyophilized proteins are shipped at ambient temperature.		