## **Nanodisc Human ACTHR Protein**



## HDFP173

## **Product Information**

<b>Product SKU</b> :	HDF	P173	Expression Host	HEK293		Size:	10µg
Target:	ACT	HR	Tag:	C-Flag Tag	l		
Additional Infor	matic	n					
<b>Conjugate</b> :		Unconjugate	d Uni	prot ID:	Q01718		
Molecular Wei	ght:	The human f	ull length ACTHR p	rotein has a l	MW of 33.9	)kDa	
Protein Informa	tion						
		NACOD					

Background:	MC2R encodes one member of the five-member G-protein associated melanocortin				
	receptor family. Melanocortins (melanocyte-stimulating hormones and				
	adrenocorticotropic hormone) are peptides derived from pro-opiomelanocortin				
	(POMC). MC2R is selectively activated by adrenocorticotropic hormone, whereas the				
	other four melanocortin receptors recognize a variety of melanocortin ligands.				
	Mutations in MC2R can result in familial glucocorticoid deficiency. Alternate				
	transcript variants have been found for this gene. [provided by RefSeq, May 2014]				
Synonyms:	ACTHR				
<b>Protein Description</b> :	Human ACTHR full length protein-synthetic nanodisc				
Formulation:	Lyophilized from nanodisc solubilization buffer (20 mM Tris-HCl, 150 mM NaCl, pH				
	8.0). Normally 5% – 8% trehalose is added as protectants before lyophilization. Please				
	8.0). Normally 5% – 8% trehalose is added as protectants before lyophilization. Please				
	8.0). Normally 5% – 8% trehalose is added as protectants before lyophilization. Please see Certificate of Analysis for specific instructions. Do not use solvents with a pH				
Protein Pathways:	8.0). Normally 5% – 8% trehalose is added as protectants before lyophilization. Please see Certificate of Analysis for specific instructions. Do not use solvents with a pH below 6.5 or those containing high concentrations of divalent metal ions (greater				
	8.0). Normally 5% – 8% trehalose is added as protectants before lyophilization. Please see Certificate of Analysis for specific instructions. Do not use solvents with a pH below 6.5 or those containing high concentrations of divalent metal ions (greater than 5 mM) in subsequent experiments.				

**Storage & Shipping**: Store at -20°C to -80°C for 12 months in lyophilized form. After reconstitution, if not intended for use within a month, aliquot and store at -80°C (Avoid repeated freezing and thawing). Lyophilized proteins are shipped at ambient temperature.