# Nanodisc Human AGRA2 Protein



## **HDFP182**

## **Product Information**

Product SKU: HDFP182 Expression Host: HEK293 Size: 10μg

**Target**: AGRA2 **Tag**: C-Flag Tag

#### **Additional Information**

**Conjugate**: Unconjugated **Uniprot ID**: Q96PE1

Molecular Weight: The human full length AGRA2 protein has a MW of 142.6kDa

#### **Protein Information**

**Background**: Endothelial receptor which functions together with RECK to enable brain endothelial

cells to selectively respond to Wnt7 signals (WNT7A or WNT7B) (PubMed:28289266,

PubMed:30026314). Plays a key role in Wnt7-specific responses, such as endothelial

cell sprouting and migration in the forebrain and neural tube, and establishment of

the blood-brain barrier (By similarity). Acts as a Wnt7-specific coactivator of canonical

Wnt signaling: required to deliver RECK-bound Wnt7 to frizzled by assembling a

higher-order RECK-ADGRA2-Fzd-LRP5-LRP6 complex (PubMed:30026314).

ADGRA2-tethering function does not rely on its G-protein coupled receptor (GPCR)

structure but instead on its combined capacity to interact with RECK extracellularly

and recruit the Dishevelled scaffolding protein intracellularly (PubMed:30026314).

Binds to the glycosaminoglycans heparin, heparin sulfate, chondroitin sulfate and

dermatan sulfate (PubMed:16982628).[UniProtKB/Swiss-Prot Function]

**Synonyms**: GPR124, TEM5

**Protein Description**: Human AGRA2 full length protein-synthetic nanodisc

Formulation: Lyophilized from nanodisc solubilization buffer (20 mM Tris-HCl, 150 mM NaCl, pH

8.0). Normally 5% – 8% trehalose is added as protectants before lyophilization. Please

see Certificate of Analysis for specific instructions. Do not use solvents with a pH

below 6.5 or those containing high concentrations of divalent metal ions (greater

than 5 mM) in subsequent experiments.

**Protein Pathways**: Angiogenesis.

**Protein Families:** Transmembrane, Druggable Genome.

**Usage**: Research use only

**Storage & Shipping**: Store at -20°C to -80°C for 12 months in lyophilized form. After reconstitution, if not

intended for use within a month, aliquot and store at -80°C (Avoid repeated freezing

and thawing). Lyophilized proteins are shipped at ambient temperature.