## **Nanodisc Human GIPR-Strep Protein**



## HDFP866

## **Product Information**

| <b>Product SKU</b> : | HDFP866          | Expression Host:      | HEK293       | Size:              | 10µg |
|----------------------|------------------|-----------------------|--------------|--------------------|------|
| Target:              | GIPR             | Tag:                  | C-Flag&St    | ер Тад             |      |
| Additional Infor     | mation           |                       |              |                    |      |
| <b>Conjugate</b> :   | Unconjugate      | ed Unip               | prot ID:     | P48546             |      |
| Molecular Wei        | ght: The human f | full length GIPR-Stre | p protein ha | s a MW of 53.2 kDa |      |
| Protein Informat     | tion             |                       |              |                    |      |

| Background:                  | A G-protein coupled receptor for gastric inhibitory polypeptide (GIP), which was           |  |  |  |
|------------------------------|--|--|--|--|
|                              | originally identified as an activity in gut extracts that inhibited gastric acid secretion |  |  |  |
|                              | and gastrin release, but subsequently was demonstrated to stimulate insulin release        |  |  |  |
|                              | in the presence of elevated glucose. Mice lacking this gene exhibit higher blood           |  |  |  |
|                              | glucose levels with impaired initial insulin response after oral glucose load. Defect in   |  |  |  |
|                              | this gene thus may contribute to the pathogenesis of diabetes.                             |  |  |  |
| Synonyms:                    | PGQTL2   |  |  |  |
| <b>Protein Description</b> : | Human GIPR-Strep full length protein-synthetic nanodisc                                    |  |  |  |
| Formulation                  | Lyophilized from nanodisc solubilization buffer (20 mM Tris-HCl, 150 mM NaCl, pH           |  |  |  |
|                              | 8.0). Normally 5% – 8% trehalose is added as protectants before lyophilization. Please     |  |  |  |
|                              | see Certificate of Analysis for specific instructions. Do not use solvents with a pH       |  |  |  |
|                              | below 6.5 or those containing high concentrations of divalent metal ions (greater          |  |  |  |
|                              | than 5 mM) in subsequent experiments.  |  |  |  |
| Protein Pathways:            | Neuroactive ligand-receptor interaction.   |  |  |  |
| Protein Families:            | Druggable Genome, GPCR, Transmembrane.   |  |  |  |
| Usage:                       | Research use only  |  |  |  |

**Storage & Shipping**: Store at -20°C to -80°C for 12 months in lyophilized form. After reconstitution, if not intended for use within a month, aliquot and store at -80°C (Avoid repeated freezing and thawing). Lyophilized proteins are shipped at ambient temperature.