## **Recombinant Human Mature MIS/AMH Protein**



## **RPCB0007**

**Product Information** 

Product SKU: RPCB0007 Gene ID: 268 Size: 10μg

**Tag**: No-Tag **Reactivity**: Human

**Additional Information** 

**Expression Host**: - **Swissprot**: P03971

**Purity**: > 95% by SDS-PAGE.

## **Protein Information**

Background:

Anti-Mullerian hormone (AMH), a member of the TGF-beta superfamily, is produced by granulosa cells (GCs) of preantral and small antral follicles and plays a role in regulating the recruitment of primordial follicles and the FSH-dependent development of follicles. BMP15 up-regulates the transcription of AMH and that the inhibition of p38 MAPK decreases the BMP15-induced expression of AMH and SOX9, suggesting that BMP15 up-regulates the expression of AMH via the p38 MAPK signaling pathway, and this process involves the SOX9 transcription factor. AMH is widely used for assessing ovarian reserve, and it is particularly convenient, because it is thought to have minimal variability throughout the menstrual cycle. Fetal anti-Mullerian hormone (AMH) is responsible for normal male sexual differentiation, and circulating AMH is used as a marker of testicular tissue in newborns with disorders of sex development. Anti-Mullerian hormone (AMH) produced in the developing testis induces the regression of the Mullerian duct, which develops into the oviducts, uterus and upper vagina. As well as other hormone receptors, and a decreased ovarian cortex cell proliferation. These results help understand the inhibitory effects of AMH on follicular development.

**Protein Description**:

High quality, high purity and low endotoxin recombinant Recombinant Human Mature MIS/AMH Protein , tested reactivity in HEK293 cells and has been validated in SDS-PAGE.100% guaranteed.

**Endotoxin**:  $< 1 \text{ EU/}\mu\text{g}$  of the protein by LAL method.

**Formulation**: Lyophilized from a 0.22 μm filtered solution of 50mM acetic acid, pH 3.0.

**Storage**: Store at -20°C. Store the lyophilized protein at -20°C to -80°C up to 1 year from the

date of receipt.After reconstitution, the protein solution is stable at -20°C for 3

months, at 2-8°C for up to 1 week.