

RPCB0375

Product Information

Product SKU:	RPCB0375	Gene ID:	93	Size:	10µg
Tag:	C-hFc&His	Reactivity:	Human		

Additional Information

Expression Host:	HEK293 cells	Swissprot:	Q13705
Purity:	> 95% by SDS-PAGE.		

Protein Information

Background: Activins are dimeric growth and differentiation factors which belong to the transforming growth factor-beta (TGF-beta) superfamily of structurally related signaling proteins. Activins signal through a heteromeric complex of receptor serine kinases which include at least two type I (I and IB) and two type II (II and IIB) receptors. These receptors are all transmembrane proteins, composed of a ligand-binding extracellular domain with cysteine-rich region, a transmembrane domain, and a cytoplasmic domain with predicted serine/threonine specificity. Type I receptors are essential for signaling; and type II receptors are required for binding ligands and for expression of type I receptors. Type I and II receptors form a stable complex after ligand binding, resulting in phosphorylation of type I receptors by type II receptors. Type II receptors are considered to be constitutively active kinases. This gene encodes activin A type IIB receptor, which displays a 3- to 4-fold higher affinity for the ligand than activin A type II receptor.

Protein Description: High quality, high purity and low endotoxin recombinant Recombinant Human Activin RII_B/ACVR2_B Protein, tested reactivity in HEK293 cells and has been validated in SDS-PAGE. 100% guaranteed.

Endotoxin: < 0.1 EU/µg of the protein by LAL method.

Formulation: Lyophilized from a 0.22 µm filtered solution of PBS, pH 7.4. Contact us for customized product form or formulation.

Contact Details | Dublin, Ireland

Email: techsupport@assaygenie.com | **Web:** www.assaygenie.com

Copyright © 2024 Assay Genie Ltd, All Rights Reserved. All information / detail is correct at time of going to print.

Storage:

Store at -20°C. Store the lyophilized protein at -20°C to -80 °C up to 1 year from the date of receipt. After reconstitution, the protein solution is stable at -20°C for 3 months, at 2-8°C for up to 1 week.