

RPCB1003

Product Information

Product SKU:	RPCB1003	Gene ID:	2056	Size:	10µg
Tag:	C-His	Reactivity:	Human		

Additional Information

Expression Host:	HEK293 cells	Swissprot:	P01588
Purity:	> 95% by SDS-PAGE.		

Protein Information

Background: Erythropoietin is a member of the EPO / TPO family. It is a secreted, glycosylated cytokine composed of four alpha helical bundles. Erythropoietin can be found in the plasma and regulates red cell production by promoting erythroid differentiation and initiating hemoglobin synthesis. It also has neuroprotective activity against a variety of potential brain injuries and antiapoptotic functions in several tissue types. Erythropoietin is the principal hormone involved in the regulation of erythrocyte differentiation and the maintenance of a physiological level of circulating erythrocyte mass. It is produced by kidney or liver of adult mammals and by liver of fetal or neonatal mammals. Genetic variation in erythropoietin is associated with susceptibility to microvascular complications of diabetes type 2. These are pathological conditions that develop in numerous tissues and organs as a consequence of diabetes mellitus. They include diabetic retinopathy, diabetic nephropathy leading to end-stage renal disease, and diabetic neuropathy. Diabetic retinopathy remains the major cause of new-onset blindness among diabetic adults. It is characterized by vascular permeability and increased tissue ischemia and angiogenesis. It has a longer circulating half-life in vivo. Erythropoietin is being much misused as a performance-enhancing drug in endurance athletes.

- Protein Description:** High quality, high purity and low endotoxin recombinant Recombinant Human Erythropoietin/EPO Protein , tested reactivity in HEK293 cells and has been validated in SDS-PAGE.100% guaranteed.
- Endotoxin:** <0.1EU/μg
- Formulation:** Lyophilized from a 0.22 μm filtered solution of PBS, pH 7.4.
- Storage:** Store at -20°C.Store the lyophilized protein at -20°C to -80 °C up to 1 year from the date of receipt.After reconstitution, the protein solution is stable at -20°C for 3 months, at 2-8°C for up to 1 week.