# **Recombinant Mouse OSMR/IL-31RB Protein**



## **RPCB1042**

### **Product Information**

**Product SKU**: RPCB1042 **Gene ID**: 18414 **Size**: 10μg

Tag: C-His Reactivity: Mouse

#### **Additional Information**

**Expression Host**: HEK293 cells **Swissprot**: O70458

**Purity**: > 95% by SDS-PAGE.

#### **Protein Information**

**Background**: Oncostatin-M specific receptor subunit beta also known as the oncostatin M receptor

(OSMR) and Interleukin-31 receptor subunit beta (IL-31RB), is one of the receptor

proteins for oncostatin M. OSMR is a member of the type I cytokine receptor family.

IL-31RB/OSMR heterodimerizes with interleukin 6 signal transducer to form the type

II oncostatin M receptor and with interleukin 31 receptor A to form the interleukin 31

receptor, and thus transduces oncostatin M and interleukin 31 induced signaling

events. Mutations in IL-31RB/OSMR have been associated with familial primary

localized cutaneous amyloidosis. Defects in IL-31RB/OSMR are the cause of

amyloidosis primary localized cutaneous type 1 (PLCA1), also known as familial lichen

amyloidosis of familial cutaneous lichen amyloidosis. PLCA1 is hereditary primary

amyloidosis characterized by localized cutaneous amyloid deposition. This condition

usually presents with itching (especially on the lower legs) and visible changes of skin

hyperpigmentation and thickening (lichenification) that may be exacerbated by

chronic scratching and rubbing. The amyloid deposits probably reflect a combination

of degenerate keratin filaments, serum amyloid P component, and deposition of

immunoglobulins.

**Protein Description**: High quality, high purity and low endotoxin recombinant Recombinant Mouse

OSMR/IL-31RB Protein, tested reactivity in HEK293 cells and has been validated in

SDS-PAGE.100% guaranteed.

**Endotoxin**: <0.1EU/μg

**Formulation**: Lyophilized from a 0.22 μm filtered solution of PBS, pH 7.4.

**Storage**: Store at -20°C.Store the lyophilized protein at -20°C to -80 °C up to 1 year from the

date of receipt. After reconstitution, the protein solution is stable at -20°C for 3  $\,$ 

months, at 2-8°C for up to 1 week.