Recombinant Human CX3CL1/Fractalkine Protein



RPCB1159

Product Information

Product SKU: RPCB1159 **Gene ID**: 6376 **Size**: 10μg

Tag: C-6His **Reactivity**: Human

Additional Information

Expression Host: Pichia **Swissprot**: P78423

Purity: -

Protein Information

Background: Fractalkine or Chemokine (C-X3-C motif) ligand 1 (CX3CL1) is a member of the CX3C

chemokine family. Fractalkine / CX3CL1 is a unique chemokine that functions not only

as a chemoattractant but also as an adhesion molecule and is expressed on endothelial cells activated by proinflammatory cytokines, such as interferon-gamma

and tumor necrosis factor-alpha. Fractalkine/CX3CL1 is expressed in a membrane-

bound form on activated endothelial cells and mediates attachment and firm

adhesion of T cells, monocytes and NK cells. Fractalkine / CX3CL1 is associated with

dendritic cells (DC) in epidermis and lymphoid organs. The fractalkine receptor,

CX3CR1, is expressed on cytotoxic effector lymphocytes, including natural killer (NK)

cells and cytotoxic T lymphocytes, which contain high levels of intracellular perforin

and granzyme B, and on macrophages. Soluble fractalkine causes migration of NK

cells, cytotoxic T lymphocytes, and macrophages, whereas the membrane-bound

form captures and enhances the subsequent migration of these cells in response to

secondary stimulation with other chemokines.

Protein Description: High quality, high purity and low endotoxin recombinant Recombinant Human

CX3CL1/Fractalkine Protein, tested reactivity in Pichia and has been validated in SDS-

PAGE.100% guaranteed.

Endotoxin: $< 0.1 EU/\mu g$

Formulation: Lyophilized from a 0.22 µm filtered solution of PBS, pH 7.4.

Storage:

Store at -20°C. Store the lyophilized protein at -20°C to -80 °C up to 1 year from the date of receipt. After reconstitution, the protein solution is stable at -20°C for 3 months, at 2-8°C for up to 1 week.