

## RPCB1219

---

**Product Information**

<b>Product SKU:</b>	RPCB1219	<b>Gene ID:</b>	16542	<b>Size:</b>	10µg
<b>Tag:</b>	C-His	<b>Reactivity:</b>	Mouse		

---

**Additional Information**

<b>Expression Host:</b>	HEK293 cells	<b>Swissprot:</b>	P35918
<b>Purity:</b>	> 90% by SDS-PAGE.		

---

**Protein Information**

**Background:** Kinase insert domain receptor (KDR) is also known as CD309, FLK1, VEGFR, VEGFR2, and is one of the subtypes of VEGFR. VEGF receptors are receptors for vascular endothelial growth factor (VEGF). VEGFR2 was shown to be the primary signal transducer for angiogenesis and the development of pathological conditions such as cancer and diabetic retinopathy. It has been shown that VEGFR2 is expressed mainly in the endothelial cells, and the expression is upregulated in the tumor vasculature. Thus the inhibition of VEGFR2 activity and its downstream signaling are important targets for the treatment of diseases involving angiogenesis. VEGFR2 transduces the major signals for angiogenesis via its strong tyrosine kinase activity. However, unlike other representative tyrosine kinase receptors, VEGFR2 does not use the Ras pathway as major downstream signaling but rather uses the phospholipase C-protein kinase C pathway to signal mitogen-activated protein (MAP)-kinase activation and DNA synthesis. VEGFR2 is a direct and major signal transducer for pathological angiogenesis, including cancer and diabetic retinopathy, in cooperation with many other signaling partners; thus, VEGFR2 and its downstream signaling appear to be critical targets for the suppression of these diseases. VEGF and VEGFR2-mediated survival signaling are critical to endothelial cell survival, maintenance of the vasculature and alveolar structure, and regeneration of lung tissue. Reduced VEGF

and VEGFR2 expression in emphysematous lungs has been linked to increased endothelial cell death and vascular regression.

- Protein Description:** High quality, high purity and low endotoxin recombinant Recombinant Mouse VEGFR-2/KDR/CD309 Protein , tested reactivity in HEK293 cells and has been validated in SDS-PAGE.100% guaranteed.
- Endotoxin:** <0.1EU/μg
- Formulation:** Lyophilized from a 0.22 μm filtered solution of PBS, pH 7.4.
- Storage:** Store at -20°C.Store the lyophilized protein at -20°C to -80 °C up to 1 year from the date of receipt.After reconstitution, the protein solution is stable at -20°C for 3 months, at 2-8°C for up to 1 week.