

RPCB1241

Product Information

Product SKU:	RPCB1241	Gene ID:	16002	Size:	10µg
Tag:	C-hFc	Reactivity:	Mouse		

Additional Information

Expression Host:	HEK293 cells	Swissprot:	P09535
Purity:	> 95% by SDS-PAGE.		

Protein Information

Background: IGF-II (Insulin-like growth factor II; also multiplication-stimulating polypeptide/MSP and somatomedin-A) is a secreted 8 kDa polypeptide that belongs to the insulin family of peptide growth factors. It is part of a complex system of growth and metabolic-regulating proteins that is particularly important during development. It has been associated with nervous system proliferation and differentiation, myelination, adrenal cortical proliferation, and skeletal growth and differentiation. In human, IGF-II is primarily synthesized by the liver, and circulates at high levels in both fetus and adult. In rodent, however, IGF-II levels drop after the perinatal period, an effect attributed to the lack of a key gene promoter. This may indicate that postnatally, IGF-II has either a limited, or local effect only in rodent. For example, evidence suggests IGF-II may be the intermediary for SHH induction of VEGF attendant with local neovascularization. Rodent cells known to express IGF-II include astrocytes, hepatocytes, osteoblasts, embryonic striated muscle cells plus Kupffer cells and Ito cells. Mouse IGF-II is synthesized as a 180 amino acid (aa) preproprecursor. It contains a 24 aa signal sequence, a 67 aa mature region, and an 89 aa C-terminal prodomain that is alternatively referred to as the E-peptide. Mature IGF-II is 91% and 97% aa identical to human and rat IGF-II, respectively. Proper processing of IGF-II requires the chaperone activity of GRP94. This generates an 8 kDa mature form, an 18 kDa, 156 aa proform, and a potential 11 kDa, 88 aa "Big"

Contact Details | Dublin, Ireland

Email: techsupport@assaygenie.com | **Web:** www.assaygenie.com

Copyright © 2024 Assay Genie Ltd, All Rights Reserved. All information / detail is correct at time of going to print.

form (aa 25-112). This 11 kDa "Big" form would be equivalent to human 15-16 kDa IGF-II, with the 5 kDa difference attributable to the presence of O-linked glycosylation . There is an additional 34 aa proteolytic fragment that is termed preptin and contains aa 93-126 of the preproprecursor. This is distinct from IGF-II, is secreted by pancreatic b cells, and facilitates insulin secretion. IGF-II has multiple binding partners. It binds to IGF-IR, the Insulin receptor (IR)-type A and IGF-IR:IR-A hybrids, the type 2 IGF receptor (IGF-2R), and IGF binding proteins 1-6 . The first three receptors initiate downstream signaling events, the IGF-2R sequesters local IGF-II, and the six IGFBPs regulate IGF-II activity in various tissues.

Protein Description: High quality, high purity and low endotoxin recombinant Recombinant Mouse IGF-II Protein , tested reactivity in HEK293 cells and has been validated in SDS-PAGE.100% guaranteed.

Endotoxin: <0.1EU/μg of the protein by LAL method.

Formulation: Lyophilized from a 0.22 μm filtered solution of PBS, pH 7.4.

Storage: Store at -20°C.Store the lyophilized protein at -20°C to -80 °C up to 1 year from the date of receipt.After reconstitution, the protein solution is stable at -20°C for 3 months, at 2-8°C for up to 1 week.