# Recombinant Human Syndecan-1/SDC1/CD138 Protein



## **RPCB1607**

## **Product Information**

**Product SKU**: RPCB1607 **Gene ID**: 6382 **Size**: 10μg

**Tag**: C-Rabbit Fc **Reactivity**: Human

### **Additional Information**

**Expression Host**: HEK293 cells **Swissprot**: P18827

**Purity**: > 95% by SDS-PAGE.

#### **Protein Information**

**Background**: Syndecan-1 also known as SDC1 and CD138, is the most extensively studied member

of the syndecan family. It is found mainly in epithelial cells, but its expression is

developmentally regulated during embryonic development. Syndecan-

1/SDC1/CD138 has been shown to mediate cell adhesion to several ECM molecules,

and to act as a coreceptor for fibroblast growth factors, potent angiogenic growth

factors involved also in differentiation. Syndecan-1/SDC1/CD138 expression is

reduced during malignant transformation of various epithelia, and this loss correlates

with the histological differentiation grade of squamous cell carcinomas, lacking from

poorly differentiated tumours. In squamous cell carcinomas of the head and neck,

positive syndecan-1 expression correlates with a more favourable prognosis.

Experimental studies on the role of Syndecan-1 in malignant transformation have

shown that Syndecan-1/SDC1/CD138 expression is associated with the maintenance

of epithelial morphology, anchorage-dependent growth and inhibition of

invasiveness in vitro.

Protein Description: High quality, high purity and low endotoxin recombinant Recombinant Human

Syndecan-1/SDC1/CD138 Protein , tested reactivity in HEK293 cells and has been

validated in SDS-PAGE.100% guaranteed.

**Endotoxin**:  $< 0.1 EU/\mu g$ 

**Formulation**: Lyophilized from a 0.22 μm filtered solution of PBS, pH 7.4.

Storage:

Store at -20°C. Store the lyophilized protein at -20°C to -80 °C up to 1 year from the date of receipt. After reconstitution, the protein solution is stable at -20°C for 3 months, at 2-8°C for up to 1 week.