

RPES8123

Product Information

Product SKU:	RPES8123	Expression Host:	E.coli	Size:	20µg
Tag:	N-Trx	Reactivity:	Human	Accession:	P02751-1

Additional Information

Calculated MW:	42.9 kDa	Observed MW:	42 kDa
Sequence:	Asp2268-Glu2477		

Protein Information

Background: Fibronectin (FN) is a glycoprotein component of the extracellular matrix of the extracellular matrix (ECM) with roles in embryogenesis, development, and wound healing. More recently, FN has emerged as player in platelet thrombus formation and diseases associated with thrombosis including vascular remodeling, atherosclerosis, and cardiac repair following a myocardial infarct. Each monomer of FN consists of three types of homologous repeating units, that is 12 type I repeats, two type II repeats and 15-17 type III repeats. The occurrence of multiple isoforms results from alternative mRNA splicing of the ED-A, ED-B and III-CS regions, and subsequent post-translational modification. As an ECM component and one of the primary cell adhesion molecules, Fibronectin can be a ligand for fibrin, heparin, chondroitin sulfate, collagen/gelatin, as well as many integrin receptors through which FN mediates the variety of cellular signaling pathways. The study of solid human tumors showed among the early signs of malignant transformation the fragmentation of pericellular FN, concomitant with the increase of its production by the peritumoral stroma. These results should encourage further investigations concerning the potential importance of Fn production and brea KDown during cancer progression. FN1 expression has been described to increase significantly from the morula towards the early blastocyst stage, suggesting that FN1 may also be involved in early blastocyst formation. The fragment 2 of FN comprises the first 7 FN type III repeats

Contact Details | Dublin, Ireland

Email: techsupport@assaygenie.com | **Web:** www.assaygenie.com

Copyright © 2024 Assay Genie Ltd, All Rights Reserved. All information / detail is correct at time of going to print.

and is suggested to be important for self association during fibril growth via the key module III2.

Synonyms:	-
Endotoxin:	< 10 EU/mg of the protein as determined by the LAL method
Formulation:	Lyophilized from a 0.2 µm filtered solution in PBS with 5% Trehalose and 5% Mannitol.
Purity:	> 90% as determined by reducing SDS-PAGE.
Bio-Activity:	Not validated for activity
Storage:	Generally, lyophilized proteins are stable for up to 12 months when stored at -20 to -80°C. Reconstituted protein solution can be stored at 4-8°C for 2-7 days. Aliquots of reconstituted samples are stable at < -20°C for 3 months.