Recombinant Human CD44 Protein (Sumo Tag)



RPES8130

Product Information

Product SKU: RPES8130 Expression Host: E.coli Size: 20μg

Tag: N-Sumo Reactivity: Human Accession: P16070-1

Additional Information

Calculated MW: 30.3 kDa Observed MW: 37 kDa

Sequence: Leu24-Ser182

Protein Information

Background: CD44 is a type I transmembrane protein and a member of the cartilage link protein

family. It is involved in cell-cell and cell-matrix interactions and signal transduction.

Several CD44 ligands have been identified. The most extensively characterized ligand

for CD44 is hyaluronan, a component of the extracellular matrix. CD44 protein is

expressed on the majority of immune cells. The binding of CD44 to hyaluronan is

induced on T lymphocytes after activation by antigen and on monocytes after

stimulation by inflammatory agents. Under inflammatory conditions, CD44 on

endothelial cells presents hyaluronan to CD44 on activated T lymphocytes and

mediates a rolling interaction under flow conditions. Perturbations of the hyaluronan-

CD44 interaction at the plasma membrane by various antagonists result in

attenuation of receptor tyrosine kinase and transporter activities and inhibition of

tumor progression in vivo. CD44 is known to interact with the ezrin family (ERM

family) members and form a complex that plays diverse roles within both normal and

abnormal cells, particularly cancer cells. CD44 and ezrin and their respective complex

have properties sµggesting that they may be important in the process of tumour-

endothelium interactions, cell migrations, cell adhesion, tumour progression and

metastasis.

Synonyms: -

Endotoxin: < 10 EU/mg of the protein as determined by the LAL method

Formulation: Lyophilized from a 0.2 μm filtered solution in PBS with 5% Trehalose and 5% Mannitol.

Purity: > 90% as determined by reducing SDS-PAGE.

Bio-Activity: Not validated for activity

Storage: Generally, lyophilized proteins are stable for up to 12 months when stored at -20 to

-80°C. Reconstituted protein solution can be stored at 4-8°C for 2-7 days. Aliquots of

reconstituted samples are stable at < -20°C for 3 months.