

RPES8282

Product Information

Product SKU:	RPES8282	Expression Host:	Mammalian	Size:	20µg
Tag:	C-Fc	Reactivity:	Human	Accession:	P41217

Additional Information

Calculated MW:	47.1 kDa	Observed MW:	60-70 kDa
Sequence:	Gln31-Gly232		

Protein Information

Background: CD200 (OX-2) is a cell surface glycoprotein that imparts immune privileges by suppressing alloimmune and autoimmune responses through its receptor, CD200R, expressed primarily on myeloid cells. Signals delivered through the CD200:CD200R axis have been shown to play an important role in the regulation of anti-tumor immunity, and overexpression of CD200 has been reported in a number of malignancies, including CLL, as well as on cancer stem cells. The role of CD200-CD200R signaling in immune regulation of the central nervous system has become a popular field of research in recent years. Many studies have shown that there is a close correlation between CD200-CD200R, microglia activation, and Parkinson's disease (PD). The ability of CD200 to suppress myeloid cell activation is critical for maintaining normal tissue homeostasis but may also enhance the survival of migratory neoplastic cells. CD200 and CD200R associate via their respective N-terminal Ig-like domains. CD200 has been characterized as an important immunoregulatory molecule, increased expression of which can lead to decreased transplant rejection, autoimmunity, and allergic disease. Elevated CD200 expression has been reported to be associated with poor prognosis in some human malignancies. Besides, CD200 also plays an important role in prevention of graft rejection, autoimmune diseases and spontaneous abortion.

Contact Details | Dublin, Ireland

Email: techsupport@assaygenie.com | **Web:** www.assaygenie.com

Copyright © 2024 Assay Genie Ltd, All Rights Reserved. All information / detail is correct at time of going to print.

Synonyms:	Antigen identified by monoclonal MRC OX 2, CD200, CD200 antigen, CD200 molecule, MOX1, MOX2, MRC, MRC OX 2 antigen, My033, OX 2, OX 2 membrane glycoprotein precursor, OX-2 membrane glycoprotein, OX2G, OX2G
Endotoxin:	< 1.0 EU/mg of the protein as determined by the LAL method
Formulation:	Lyophilized from a 0.2 µm filtered solution in PBS with 5% Trehalose and 5% Mannitol.
Purity:	> 90% as determined by reducing SDS-PAGE.
Bio-Activity:	Not validated for activity
Storage:	Generally, lyophilized proteins are stable for up to 12 months when stored at -20 to -80°C. Reconstituted protein solution can be stored at 4-8°C for 2-7 days. Aliquots of reconstituted samples are stable at < -20°C for 3 months.