Recombinant human CDNF Protein (Fc Tag)



RPES8312

Product Information

Product SKU: Tag:	RPES8312 C-Fc	Expression Host: Reactivity:	Mammalian human	Size: Accession:	20µg Q49АН0	
Additional Information						
Calculated MW	/: 45.4 kDa	Obse	rved MW:	40-50 kDa		
Sequence:	Met1-Leu187	7				

Protein Information

Background :	Cerebral Dopamine Neurotrophic Factor (CDNF), also known as ARMETL1 (ARMET-		
	like protein 1), is a secreted protein with eight conserved cysteine residues, predicting		
	a unique protein fold and defining a new, evolutionarily conserved protein family.		
	CDNF is a novel neurotrophic factor with strong trophic activity on dopaminergic		
	neurons comparable to that of glial cell line-derived neurotrophic factor (GDNF).		
	CDNF/ARMETL1 is a evolutionary conserved protein which can protect and restore		
	the function of dopaminergic neurons in the rat model of Parkinson's disease,		
	suggesting that CDNF might be beneficial for the treatment of Parkinson's disease.		
	CDNF is widely expressed in neurons in several brain regions including cerebral		
	cortex, hippocampus, substantia nigra, striatum and cerebellum. Human CDNF is		
	glycosylated and secreted from transiently transfected cells. CDNF promotes the		
	survival, growth, and function of dopamine-specific neurons and is expressed in brain		
	regions that undergo cocaine-induced neuroplasticity.		
Synonyms:	Arginine rich mutated in early stage tumors like 1, ARMET L1, ARMET like protein 1,		
	ARMET-like protein 1, ARMETL 1, CDNF, CDNF, Cerebral dopamine neurotrophic		
	factor, Conserved dopamine neurotrophic factor		
Endotoxin :	< 1.0 EU/mg of the protein as determined by the LAL method		
Formulation	Lyophilized from a 0.2 μm filtered solution in PBS with 5% Trehalose and 5% Mannitol.		
Purity:	> 90% as determined by reducing SDS-PAGE.		

Bio-Activity:Not validated for activityStorage:Generally, lyophilized proteins are stable for up to 12 months when stored at -20 to
-80°C. Reconstituted protein solution can be stored at 4-8°C for 2-7 days. Aliquots of
reconstituted samples are stable at < -20°C for 3 months.</th>