Recombina Tag) RPES8334					
Product Informa	ation				
Product SKU:	RPES8334	Expression Host:	Mammalian	Size:	20µg
Tag:	C-Fc	Reactivity:	Human	Accession :	Q9P0V8
dditional Inform	mation				
Calculated MW	: 48.1 kDa	Obse	erved MW: 55 kDa		
Sequence:	Ala23-Asp23	3			

۸

Protein Information

- Background: The signaling lymphocyte activation molecule (SLAM) family includes homophilic and heterophilic receptors that modulate both adaptive and innate immune responses. These receptors share a common ectodomain organization: a membrane-proximal immunoglobulin constant domain and a membrane-distal immunoglobulin variable domain that is responsible for ligand recognition. SLAM family of receptors is expressed by a wide range of immune cells. Through their cytoplasmic domain, SLAM family receptors associate with SLAM-associated protein (SAP)-related molecules , a group of cytoplasmic adaptors composed almost exclusively of an SRC homology 2 domain. SLAM family receptors, in association with SAP family adaptors, have crucial roles during normal immune reactions in innate and adaptive immune cells. Mouse SLAM family member 8, also known as B-lymphocyte activator macrophage expressed, BCM-like membrane protein, SLAMF8 and BLAME, is a single-pass type I membrane protein. It contains one Ig-like C2-type (immunoglobulin-like) domain. SLAMF8 / BLAME is expressed in lymph node, spleen, thymus and bone marrow. It may play a role in B-lineage commitment and/or modulation of signaling through the B-cell receptor. Synonyms: SLAM family member, SBBI, SLAF, Slamf, BCM-like membrane protein, B-lymphocyte
 - SLAM family member, SBBI, SLAF, Slamf, BCM-like membrane protein, B-lymphocyte activator macrophage expressed, CD353, SBBI42, SLAF8, SLAM family member 8, BLAME, SLAMF8, BLAME, SBBI42

Endotoxin:	< 1.0 EU/mg of the protein as determined by the LAL method		
Formulation :	Lyophilized from a 0.2 μm filtered solution in PBS with 5% Trehalose and 5% Mannitol.		
Purity:	> 90% as determined by reducing SDS-PAGE.		
Bio-Activity :	Not validated for activity		
Storage:	Generally, lyophilized proteins are stable for up to 12 months when stored at -20 to		
	-80°C. Reconstituted protein solution can be stored at 4-8°C for 2-7 days. Aliquots of		
	reconstituted samples are stable at $< -20^{\circ}$ C for 3 months.		