

RPES8394

Product Information

Product SKU:	RPES8394	Expression Host:	Mammalian	Size:	20µg
Tag:	C-Fc	Reactivity:	Mouse	Accession:	Q9R182

Additional Information

Calculated MW:	73.2 kDa	Observed MW:	90-100 kDa
Sequence:	Ser17-Thr455		

Protein Information

Background: ANGPTL3 is a secreted glycoprotein that is structurally related to the angiopoietins. Mature mouse ANGPTL3 contains an N-terminal coiled coil domain and a C-terminal fibrinogen-like domain. ANGPTL3 is expressed in the liver from early in development through adulthood. Full length ANGPTL3 circulates in the plasma as do the proteolytically separated N- and C-terminal fragments containing the coiled-coil domain and fibrinogen-like domains, respectively. ANGPTL3 is found as 70 kDa, 50 kDa, and 32 kDa species and can form weakly associated noncovalent multimers in vitro. ANGPTL3 directly inhibits lipoprotein lipase (LPL), an enzyme responsible for hydrolyzing circulating triglycerides. This activity requires a putative heparin binding motif that is N-terminal to the coiled coil domain. Proteolytic removal of the fibrinogen-like domain from the N-terminal fragment serves to activate ANGPTL3 and increase its ability to inhibit LPL in vitro and function in vivo. ANGPTL3 promotes an increase in circulating triglyceride levels without altering VLDL or HDL secretion or uptake. ANGPTL3 knockout mice are hypolipidemic and have elevated LPL activity. ANGPTL3 expression in vivo is up regulated by LXR agonists and down regulated by insulin, leptin, and TR beta agonists. Dysregulated ANGPTL3 expression and elevated plasma triglyceride levels are characteristic of some strains of obese and diabetic mice. ANGPTL3 does not bind Tie-1 or Tie-2 but its fibrinogen-like domain interacts with integrin alpha V beta 3 to induce endothelial cell adhesion, migration, and

Contact Details | Dublin, Ireland

Email: techsupport@assaygenie.com | **Web:** www.assaygenie.com

Copyright © 2024 Assay Genie Ltd, All Rights Reserved. All information / detail is correct at time of going to print.

neovascularization. ANGPTL3, secreted by fetal liver cells, also promotes the expansion of hematopoietic stem cells. Mature mouse ANGPTL3 shares 22%-30% amino acid (aa) sequence identity with ANGPTL1, 2, 4, 6, and 7. It shares 77% aa sequence identity with human ANGPTL3.

Synonyms:	-
Endotoxin:	< 1.0 EU/mg of the protein as determined by the LAL method
Formulation:	Lyophilized from a 0.2 µm filtered solution in PBS with 5% Trehalose and 5% Mannitol.
Purity:	> 90% as determined by reducing SDS-PAGE.
Bio-Activity:	Not validated for activity
Storage:	Generally, lyophilized proteins are stable for up to 12 months when stored at -20 to -80°C. Reconstituted protein solution can be stored at 4-8°C for 2-7 days. Aliquots of reconstituted samples are stable at < -20°C for 3 months.