

RPES8418

Product Information

Product SKU: RPES8418	Expression Host: Mammalian	Size: 20µg
Tag: C-Fc	Reactivity: Mouse	Accession: P20334

Additional Information

Calculated MW: 42.93 kDa	Observed MW: 50-60 kDa
Sequence: Val24-Leu187	

Protein Information

Background: CD137 (also known as 4-1BB) is a surface co-stimulatory glycoprotein originally described as present on activated T lymphocytes, which belongs to the tumor necrosis factor (TNF) receptor superfamily. It is expressed mainly on activated CD4+ and CD8+ T cells, and binds to a high-affinity ligand (4-1BBL) expressed on several antigen-presenting cells such as macrophages and activated B cells. Upon ligand binding, 4-1BB is associated with the tumor necrosis factor receptor-associated factors (TRAFs), the adaptor protein which mediates downstream signaling events including the activation of NF-kappaB and cytokine production. 4-1BB signaling either by binding to 4-1BBL or by antibody ligation delivers signals for T-cell activation and growth, as well as monocyte proliferation and B-cell survival, and plays an important role in the amplification of T cell-mediated immune responses. In addition, CD137 and CD137L are expressed in different human primary tumor tissues, suggesting that they may influence the progression of tumors. Crosslinking of CD137 on activated T cells has shown promise in enhancing anti-tumor immune responses in murine models, and agonistic anti-CD137 antibodies are currently being tested in phase I clinical trials. Soluble forms of CD137 (sCD137) are generated by differential splicing. sCD137 can bind to CD137 ligand to antagonize the costimulatory activities of the membrane-bound CD137 and reduce T cell proliferation and IL-2 secretion.

Contact Details | Dublin, Ireland

Email: techsupport@assaygenie.com | **Web:** www.assaygenie.com

Copyright © 2024 Assay Genie Ltd, All Rights Reserved. All information / detail is correct at time of going to print.

Synonyms:	4 1BB, 4 1BB ligand receptor, 4-1BB ligand receptor, 4-1BB Ligand Receptor T Cell, 4-1BB, mouse, homolog of, Antigen 4-1BB Homolog, CD 137, CD137, CD137 antigen, CDw137, HLDA VI, Homolog of mouse 4 1BB, ILA, induced by lymphocyte activation, ILA, Induced by lymphocyte activation, Interleukin activated receptor homolog of mouse Ly63, Ly63, mouse, homolog of, MGC2172, OTTHUMP00000044294, Receptor protein 4 1BB, T cell antigen 4 1BB homolog, T cell antigen ILA, T-cell antigen 4-1BB homolog, T-cell antigen ILA, TNF receptor superfamily member 9, TNFRSF9, TNFR9, Tumor necrosis factor receptor superfamily member 9
Endotoxin:	< 1.0 EU/mg of the protein as determined by the LAL method
Formulation:	Lyophilized from a 0.2 µm filtered solution in PBS with 5% Trehalose and 5% Mannitol.
Purity:	> 90% as determined by reducing SDS-PAGE.
Bio-Activity:	Not validated for activity
Storage:	Generally, lyophilized proteins are stable for up to 12 months when stored at -20 to -80°C. Reconstituted protein solution can be stored at 4-8°C for 2-7 days. Aliquots of reconstituted samples are stable at < -20°C for 3 months.