

RPES8435

Product Information

Product SKU: Tag:	RPES8435 C-His	Expression Host: Reactivity:	Mammalia Mouse	n Size: Accession:	20µg Q00609-1	
Additional Information						
Calculated MW	!: 22.8 kDa	Obse	erved MW:	40-60 kDa		
Sequence:	Val38-Asn24	6				

Protein Information

Background:	The B-lymphocyte activation antigen B7-1 (referred to as B7), also known as CD80, is		
	a member of cell surface immunoglobulin superfamily and is expressed on the		
	surface of antigen-presenting cells including activated B cells, macrophages and		
	dendritic cells. As costimulatory ligands, B7-1 which exists predominantly as dimer		
	and the related protein B7-2, interact with an the costimulatory receptors CD28 and		
	cytotoxic T lymphocyte-associated antigen 4 (CTLA-4) expressed on T cells, and thus		
	constitute one of the dominant pathways that regulate T cell activation and tolerance,		
	cytokine production, and the generation of CTL. The B7/CD28/CTLA4 pathway has		
	the ability to both positively and negatively regulate immune responses. CD80 is thus		
	regarded as promising therapeutic targets for autoimmune diseases and various		
	carcinomas.		
Synonyms:	T-lymphocyte activation antigen CD80, Cd80, Activation B7-1 antigen, B7,		
Endotoxin:	< 1.0 EU/mg of the protein as determined by the LAL method		
Formulation:	Lyophilized from a 0.2 μm filtered solution in PBS with 5% Trehalose and 5% Mannitol.		
Purity:	> 90% as determined by reducing SDS-PAGE.		
Bio-Activity :	Not validated for activity		
Storage:	Generally, lyophilized proteins are stable for up to 12 months when stored at -20 to		
	-80°C. Reconstituted protein solution can be stored at 4-8°C for 2-7 days. Aliquots of		
	reconstituted samples are stable at $< -20^{\circ}$ C for 3 months.		

Contact Details | Dublin, Ireland Email: techsupport@assaygenie.com | Web: <u>www.assaygenie.com</u> Copyright © 2024 Assay Genie Ltd, All Rights Reserved. All information / detail is correct at time of going to print.