

Recombinant Mouse Folate Binding protein/FBP Protein (Fc Tag)

RPES8449



Product Information

Product SKU:	RPES8449	Expression Host:	Mammalian	Size:	20µg
Tag:	C-Fc	Reactivity:	Mouse	Accession:	P35846

Additional Information

Calculated MW:	47.8 kDa	Observed MW:	55-60 kDa
Sequence:	Thr25-Ser232		

Protein Information

Background: The protein encoded by FOLR1 gene is a member of the folate receptor family. Members of this gene family bind folic acid and its reduced derivatives, and transport 5-methyltetrahydrofolate into cells. This gene product is a secreted protein that either anchors to membranes via a glycosyl-phosphatidylinositol linkage or exists in a soluble form. Mutations in this gene have been associated with neurodegeneration due to cerebral folate transport deficiency. Due to the presence of two promoters, multiple transcription start sites, and alternative splicing, multiple transcript variants encoding the same protein have been found for this gene. Folate receptor α (FR α) is the most important subunit of Folate receptor and the alpha isoform has been shown to be selectively overexpressed in cancer types like breast and ovarian cancer compared to normal breast and ovarian epithelial cells. It was determined that Folate receptor α exhibits a limited expression on the apical surfaces of the epithelial cells of normal lung, breast, thyroid, parathyroid, and kidney tissues. For their uptake of folate, normal cells rely almost exclusively on the reduced folate carrier, whereas many carcinomas and myeloid leukemia cells overexpress a high-affinity FR on their surfaces, perhaps reflecting their increased need for folate to support rapid cell division.

Synonyms: adult, Adult folate binding protein, Adult folate-binding protein, FBP, Folate Binding Protein, Folate receptor, Folate receptor 1, Folate Receptor 1 Adult, Folate Receptor

Contact Details | Dublin, Ireland

Email: techsupport@assaygenie.com | Web: www.assaygenie.com

Copyright © 2024 Assay Genie Ltd, All Rights Reserved. All information / detail is correct at time of going to print.

1 Precursor, Folate receptor adult, Folate receptor alpha, FOLR, FOLR1, FOLR1_HUMAN, FR alpha, FR-alpha, FRalpha KB cells FBP MOV18, Ovarian cancer associated antigen, Ovarian tumor associated antigen, Ovarian tumor associated antigen MOV18, Ovarian tumor-associated antigen MOV18

Endotoxin: < 1.0 EU/mg of the protein as determined by the LAL method

Formulation: Lyophilized from a 0.2 µm filtered solution in PBS with 5% Trehalose and 5% Mannitol.

Purity: > 90% as determined by reducing SDS-PAGE.

Bio-Activity: Not validated for activity

Storage: Generally, lyophilized proteins are stable for up to 12 months when stored at -20 to -80°C. Reconstituted protein solution can be stored at 4-8°C for 2-7 days. Aliquots of reconstituted samples are stable at < -20°C for 3 months.