Recombinant Mouse CD6 Protein (His Tag)



RPES8451

Product Information

Product SKU: RPES8451 Expression Host: Mammalian Size: 20μg

Tag: C-His Reactivity: Mouse Accession: Q61003

Additional Information

Calculated MW: 41.9 kDa Observed MW: 60-80 kDa

Sequence: Gly17-Thr398

Protein Information

Background: T-cell differentiation antigen CD6, also known as TP12 and CD6, is a single-pass type

I membrane protein which contains threeSRCR domains. CD6 / TP12 is a cell surface

glycoprotein expressed primarily on T cells, it may function as a costimulatory

molecule and may play a role in autoreactive immune responses. CD6 / TP12 is expressed by thymocytes, mature T-cells, a subset of B-cells known as B-1 cells, and

by some cells in the brain. CD6 ligand termed CD166 (previously known as activated

leukocyte cell adhesion molecule, ALCAM) has been identified and shown to be

expressed on activated T cells, B cells, thymic epithelium, keratinocytes, and in

rheumatoid arthritis synovial tissue. CD6 / TP12 binds to activated leukocyte cell

adhesion molecule (CD166), and is considered as a costimulatory molecule involved

in lymphocyte activation and thymocyte development. CD6 / TP12 partially associates

with the TCR / CD3 complex and colocalizes with it at the center of the mature $\ensuremath{\mathsf{CD}}$

immunological synapse (IS) on T lymphocytes. During thymic development CD6-

dependent signals may contribute both to thymocyte survival, and to the overall

functional avidity of selection in both man and mouse.

Synonyms: CD_antigen=CD6, CD6, CD6 antigen Tp120, CD6 molecule, CD6, FLJ44171, OX52, T

cell differentiation antigen CD6, T-cell differentiation antigen CD6, T12, TP120

Endotoxin: < 1.0 EU/mg of the protein as determined by the LAL method

Formulation: Lyophilized from a 0.2 μm filtered solution in PBS with 5% Trehalose and 5% Mannitol.

Purity: > 90% as determined by reducing SDS-PAGE.

Bio-Activity: Not validated for activity

Storage: Generally, lyophilized proteins are stable for up to 12 months when stored at -20 to

-80°C. Reconstituted protein solution can be stored at 4-8°C for 2-7 days. Aliquots of

reconstituted samples are stable at < -20°C for 3 months.